

UNIT – I: RELATIONS AND FUNCTIONS

CHAPTER-1

RELATIONS AND FUNCTIONS

Topic-1

Relations

Concepts Covered • Types of relations and their identification • Equivalence class



Revision Notes

1. Definition

A relation R , from a non-empty set A to another non-empty set B is mathematically as an subset of $A \times B$. Equivalently, any subset of $A \times B$ is a relation from A to B .

Thus, R is a relation from A to B

$$\Leftrightarrow R \subseteq A \times B$$

$$\Leftrightarrow R \subseteq \{(a, b) : a \in A, b \in B\}$$

Note:

- Let A and B be two non-empty finite sets having p and q elements respectively. Then $n(A \times B) = n(A) \cdot n(B) = pq$. Then total number of subsets of $A \times B = 2^{pq}$. Since each subset of $A \times B$ is a relation from A to B , therefore total number of relations from A to B will be 2^{pq} .

2. Domain & Range of a Relation

(a) **Domain of a Relation:** Let R be a relation from A to B . The domain of relation R is the set of all those elements $a \in A$ such that $(a, b) \in R \forall b \in B$. Thus, $\text{Dom.}(R) = \{a \in A : (a, b) \in R \forall b \in B\}$.

That is, the domain of R is the set of first components of all the ordered pairs which belong to R .

(b) **Range of a Relation:** Let R be a relation from A to B . The range of relation R is the set of all those elements $b \in B$ such that $(a, b) \in R \forall a \in A$.

Thus, $\text{Range of } R = \{b \in B : (a, b) \in R \forall a \in A\}$.

That is, the range of R is the set of second components of all the ordered pairs which belong to R .

(c) **Co-domain of a Relation:** Let R be a relation from A to B . Then B is called the co-domain of the relation R . So we can observe that co-domain of a relation R from A into B is the set B as a whole.

For example, Let $a \in A$ and $b \in B$ and

- (i) Let $A = \{1, 2, 3, 7\}$,
 $B = \{3, 6\}$. If aRb means $a < b$.

Then we have

$$R = \{(1, 3), (1, 6), (2, 3), (2, 6), (3, 6)\}.$$

Here, $\text{Dom.}(R) = \{1, 2, 3\}$,

$\text{Range of } R = \{3, 6\}$, $\text{Co-domain of } R = B = \{3, 6\}$

3. Types of relations from one set to another set

(a) **Empty relation:** A relation R from A to B is called an empty relation or a void relation from A to B if $R = \phi$.

For example, Let

$$A = \{2, 4, 6\}, B = \{7, 11\}$$

Let $R = \{(a, b) : a \in A, b \in B \text{ and } |a-b| \text{ is even}\}$.

Here R is an empty relation.

(b) **Universal relation:** A relation R from A to B is said to be the universal relation if $R = A \times B$.

For example, Let

$$A = \{1, 2\}, B = \{1, 3\}$$

Let $R = \{(1, 1), (1, 3), (2, 1), (2, 3)\}$.

Here, $R = A \times B$, so relation R is a universal relation.

Note:

- The void relations i.e., ϕ and universal relation are respectively the smallest and largest relations defined on the set A . Also these are also called **Trivial Relations** and other relation is called a **Non-Trivial Relation**.
- The relations $R = \phi$ and $R = A \times A$ are two **extreme relations**.

(c) **Identity relation:** A relation R defined on a set A is said to be the identity relation on A if

$$R = \{(a, b) : a \in A, b \in A \text{ and } a = b\}$$

Thus identity relation

$$R = \{(a, a) : \forall a \in A\}$$

The identity relation on set A is also denoted by I_A .

For example, Let $A = \{1, 2, 3, 4\}$,

$$\text{Then } I_A = \{(1, 1), (2, 2), (3, 3), (4, 4)\}$$

But the relation given by

$$R = \{(1, 1), (2, 2), (1, 3), (4, 4)\}$$

is not an identity relation because element of I_A is not related to elements 1 and 3.

(d) **Reflexive relation:** A relation R defined on a set A is said to be reflexive if $a R a \forall a \in A$ i.e., $(a, a) \in R \forall a \in A$.

For example, Let $A = \{1, 2, 3\}$ and R_1, R_2, R_3 be the relations given as

$$R_1 = \{(1, 1), (2, 2), (3, 3)\},$$

$$R_2 = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 1), (1, 3)\} \text{ and}$$

$$R_3 = \{(2, 2), (2, 3), (3, 2), (1, 1)\}$$

Here R_1 and R_2 are reflexive relations on A but R_3 is not reflexive as $3 \in A$ but $(3, 3) \notin R_3$.

Note:

- The identity relation is always a reflexive relation but the converse may or may not be true. As shown in the example above, R_1 is both identity as well as reflexive relation on A but R_2 is only reflexive relation on A .

(e) **Symmetric relation:** A relation R defined on a set A is symmetric if

$$(a, b) \in R \Rightarrow (b, a) \in R \quad \forall a, b \in A \text{ i.e., } aRb \Rightarrow bRa$$

(i.e., whenever aRb then bRa).

For example, Let $A = \{1, 2, 3\}$,

$$R_1 = \{(1, 2), (2, 1)\}, R_2 = \{(1, 2), (2, 1), (1, 3), (3, 1)\}.$$

$$R_3 = \{(1, 2), (2, 1), (1, 3), (3, 1), (2, 3), (3, 2)\}$$

$$R_4 = \{(1, 3), (3, 1), (2, 3)\}$$

Here R_1, R_2 and R_3 are symmetric relations on A . But R_4 is not symmetric because $(2, 3) \in R_4$ but $(3, 2) \notin R_4$.

(f) **Transitive relation:** A relation R on a set A is transitive if $(a, b) \in R$ and $(b, c) \in R \Rightarrow (a, c) \in R$ i.e., aRb and $bRc \Rightarrow aRc$.

For example, Let $A = \{1, 2, 3\}$,

$$R_1 = \{(1, 2), (2, 3), (1, 3), (3, 2)\}$$

and $R_2 = \{(1, 3), (3, 2), (1, 2)\}$

Here R_2 is transitive relation whereas R_1 is not transitive because $(2, 3) \in R_1$ and $(3, 2) \in R_1$ but $(2, 2) \notin R_1$.

(g) **Equivalence relation:** Let A be a non-empty set,

then a relation R on A is said to be an equivalence relation if

- R is reflexive.
- R is symmetric.
- R is transitive.

For example, Let $A = \{1, 2, 3\}$

$$R = \{(1, 2), (1, 1), (2, 1), (2, 2), (3, 3), (1, 3), (3, 1), (3, 2), (2, 3)\}$$

Here R is reflexive, symmetric and transitive. So R is an equivalence relation on A .

Equivalence classes: Let A be an equivalence relation in a set A and let $a \in A$. Then, the set of all those elements of A which are related to a , is called equivalence class determined by a and it is denoted by $[a]$. Thus, $[a] = \{b \in A : (a, b) \in A\}$

**Mnemonics**

Concept: Types of relation

Mnemonics: RIPE STRAWBERRY TO EAT

Interpretations

Ripe	–	Reflexive
Strawberry	–	Symmetric
To	–	Transitive
Eat	–	Equivalence

4. Inverse relation

Let $R \subseteq A \times B$ be a relation from A to B . Then, the inverse relation of R , to be denoted by R^{-1} , is a relation from B to A defined by $R^{-1} = \{(b, a) : (a, b) \in R\}$

Thus $(a, b) \in R \Leftrightarrow (b, a) \in R^{-1} \quad \forall a \in A, b \in B$.

Clearly, **Domain (R^{-1}) = Range of R ,**

Range of R^{-1} = Domain (R).

Also, $(R^{-1})^{-1} = R$.

**Key Facts**

- A relation R from A to B is an empty relation or void relation if $R = \phi$
 - A relation R on a set A is an empty relation or void relation if $R = \phi$
- A relation R from A to B is a universal relation if $R = A \times B$.
 - A relation R on a set A is an universal relation if $R = A \times A$.
- A relation R on a set A is reflexive if $aRa, \forall a \in A$.
- A relation R on a set A is symmetric if whenever aRb , then bRa for all $a, b \in A$.
- A relation R on a set A is transitive if whenever aRb and bRc then aRc for all $a, b, c \in A$.
- A relation R on A is identity relation if $R = \{(a, a) \mid a \in A\}$ i.e., R contains only elements of the type $(a, a) \forall a \in A$ and it contains no other element.

Example 1

Let N denote the set of all natural numbers and R be the relation on $N \times N$ defined by $(a, b) R(c, d)$ if $ad(b + c) = bc(a + d)$. Show that R is an equivalence relation.

Sol.

Step I : Given $(a, b) R(c, d)$ as $ad(b + c) = bc(a + d)$

$$\therefore \forall a, b \in N$$

$$\text{or } ab(b + a) = ba(a + b)$$

$$\text{or } (a, b) R(a, b)$$

$\therefore R$ is reflexive. ...(i)

Step II : Let $(a, b) R(c, d)$ for $(a, b), (c, d) \in N \times N$

$$\therefore ad(b + c) = bc(a + d) \quad \dots\text{(ii)}$$

$$\text{Also, } (c, d) R(a, b)$$

$$\therefore cb(d + a) = da(c + b)$$

[By commutation of addition and multiplication on N]

$\therefore R$ is symmetric. ...(iii)

Step III : Let $(a, b) R(c, d)$ and $(c, d) R(e, f)$ for $a, b, c, d, e, f \in N$

$$\therefore ad(b + c) = bc(a + d) \quad \dots\text{(iv)}$$

$$\text{and } cf(d + e) = de(c + f) \quad \dots\text{(v)}$$

Dividing eqn. (iv) by $abcd$ and eqn. (v) by $cdef$

$$\text{i.e., } \frac{1}{c} + \frac{1}{b} = \frac{1}{d} + \frac{1}{a}$$

$$\text{and } \frac{1}{e} + \frac{1}{d} = \frac{1}{f} + \frac{1}{c}$$

On adding, we get

$$\frac{1}{c} + \frac{1}{b} + \frac{1}{e} + \frac{1}{d} = \frac{1}{d} + \frac{1}{a} + \frac{1}{f} + \frac{1}{c}$$

$$\text{or } af(b + e) = be(a + f)$$

Hence, $(a, b) R(e, f)$

$\therefore R$ is transitive. ...(vi)

From equations (i), (iii) and (vi), R is an equivalence relation.

Topic-2**Functions**

Concepts Covered • Types of functions and their identification

**Revision Notes**

- Function as a special type of relation:** A relation f from a set A to another set B is said to be a function (or mapping) from A to B if with every element (say x) of A , the relation f relates a unique element (say y) of B . This y is called f -image of x . Also x is called pre-image of y under f .
- Difference between relation and function:** A relation from a set A to another set B is any subset of $A \times B$; while a function f from A to B is a subset of $A \times B$ satisfying following conditions:
 - For every $x \in A$, there exists $y \in B$ such that $(x, y) \in f$.
 - If $(x, y) \in f$ and $(x, z) \in f$ then, $y = z$.

S. No.	Function	Relation
(i)	Each element of A must be related to some element of B .	There may be some elements of A which are not related to any element of B .
(ii)	An element of A should not be related to more than one element of B .	An element of A may be related to more than one element of B .

- Real valued function of a real variable:** If the domain and range of a function f are subsets of R (the set of real numbers), then f is said to be a **real valued function of a real variable** or a **real function**.

4. Some important real functions and their domain & range

S. No.	Function	Representation	Domain	Range
(i)	Identity function	$I(x) = x \forall x \in \mathbb{R}$	\mathbb{R}	\mathbb{R}
(ii)	Modulus function or Absolute value function	$f(x) = x = \begin{cases} -x, & \text{if } x < 0 \\ x, & \text{if } x \geq 0 \end{cases}$	\mathbb{R}	$[0, \infty)$

S. No.	Function	Representation	Domain	Range
(iii)	Greatest integer function or Integral function or Step function	$f(x) = [x] \forall x \in \mathbb{R}$	\mathbb{R}	\mathbb{Z}
(iv)	Signum function	$f(x) = \begin{cases} x & \text{if } x \neq 0 \\ x & \text{if } x = 0 \end{cases}$ i.e., $f(x) = \begin{cases} 1, & x > 0 \\ 0, & x = 0 \\ -1, & x < 0 \end{cases}$	\mathbb{R}	$\{-1, 0, 1\}$
(v)	Exponential function	$f(x) = a^x, \forall a > 0, a \neq 1$	\mathbb{R}	$(0, \infty)$
(vi)	Logarithmic function	$f(x) = \log_a x, \forall a \neq 1, a > 0 \text{ and } x > 0$	$(0, \infty)$	\mathbb{R}

5. Types of Function

(a) **One-one function (Injective function or Injection):** A function $f : A \rightarrow B$ is one-one function or injective function if distinct elements of A have distinct images in B .

Thus, $f : A \rightarrow B$ is one-one $\Leftrightarrow f(a) = f(b)$

$$\Rightarrow a = b, \forall a, b \in A$$

$$\Leftrightarrow a \neq b \Rightarrow f(a) \neq f(b) \forall a, b \in A.$$

- If A and B are two sets having m and n elements respectively such that $m \leq n$, then total number of one-one functions from set A to set B is ${}^n C_m \times m!$ i.e., ${}^n P_m$.
- If $n(A) = n$, then the number of injective functions defined from A onto itself is $n!$.

ALGORITHM TO CHECK THE INJECTIVITY OF A FUNCTION

STEP 1: Take any two arbitrary elements a, b in the domain of f .

STEP 2: Put $f(a) = f(b)$.

STEP 3: Solve $f(a) = f(b)$. If it gives $a = b$ only, then f is a one-one function.

(b) **Onto function (Surjective function or Surjection):** A function $f : A \rightarrow B$ is onto function or a surjective function if every element of B is the f -image of some element of A . That implies $f(A) = B$ or range of f is the co-domain of f .

Thus, $f : A \rightarrow B$ is onto $\Leftrightarrow f(A) = B$ i.e., range of $f =$ co-domain of f .

ALGORITHM TO CHECK THE SURJECTIVITY OF A FUNCTION

STEP 1: Take an element $b \in B$, where B is the co-domain of the function.

STEP 2: Put $f(x) = b$.

STEP 3: Solve the equation $f(x) = b$ for x and obtain x in terms of b . Let $x = g(b)$.

STEP 4: If for all values of $b \in B$, the values of x obtained from $x = g(b)$ are in A , then f is onto. If there are some $b \in B$ for which values of x , given

by $x = g(b)$, is not in A . Then f is not onto.



Mnemonics

Concept: Types of functions

Mnemonics: Indian Syndicate Bank

Interpretations

Indian	–	injective
Syndicate	–	surjective
Bank	–	Bijjective

Also note that a bijective function is also called a one-to-one function or one-to-one correspondence.

If $f : A \rightarrow B$ is a function such that,

- f is one-one $\Rightarrow n(A) \leq n(B)$.
- f is onto $\Rightarrow n(B) \leq n(A)$.

For an ordinary finite set A , a one-one function $f : A \rightarrow A$ is necessarily onto and an onto function $f : A \rightarrow A$ is necessarily one-one for every finite set A .

(d) **Identity function:** The function $I_A : A \rightarrow A; I_A(x) = x, \forall x \in A$ is called an identity function on A .

Note:

- Domain $(I_A) = A$ and Range $(I_A) = A$.

(e) **Equal function:** Two functions f and g having the same domain D are said to be equal if $f(x) = g(x)$ for all $x \in D$.

6. Defining a Function

Consider A and B be two non-empty sets, then a rule f which associates **each element of A with a unique element of B** is called a function or the mapping from A to B or f maps A to B . If f is a mapping from A to B , then we write $f : A \rightarrow B$ which is read as ' f is mapping from A to B ' or ' f is a function from A to B '.

If f associates $a \in A$ to $b \in B$, then we say that ' b is the **image of the element a under the function f** ' or ' b is the **f -image of a** ' or '**the value of f at a** ' and denotes it by $f(a)$ and we write $b = f(a)$. The element a is called the **pre-image or inverse-image of b** .

Thus for a bijective function from A to B ,

- A and B should be non-empty.
- Each element of A should have image in B .

- (c) No element of A should have more than one image in B .
- (d) If A and B have respectively m and n number of elements then the **number of functions defined from A to B is n^m** .

7. Domain, Co-domain and Range of A function

The set A is called the **domain** of the function f and the set B is called the **co-domain**. The set of the images of

all the elements of A under the function f is called the **range of the function f** and is denoted as $f(A)$.

Thus range of the function f is $f(A) = \{f(x) : x \in A\}$.

Clearly $f(A) = B$ for a bijective function.

Example 1

Determine whether the function $f: A \rightarrow B$ defined by $f(x) = 4x + 7, x \in A$ is **one-one**.

Show that no two elements in domain have same image in codomain.

Solution :

Given, $f: A \rightarrow B$ defined by $f(x) = 4x + 7, x \in A$

Let, $x_1, x_2 \in A$, such that $f(x_1) = f(x_2)$

$$\Rightarrow 4x_1 + 7 = 4x_2 + 7 \Rightarrow 4x_1 = 4x_2 \Rightarrow x_1 = x_2$$

So, f is one-one function.

CHAPTER-2

INVERSE TRIGONOMETRIC FUNCTIONS



Revision Notes

As we have learnt in class XI, the domain and range of trigonometric functions are given below:

S. No.	Function	Domain	Range
(i)	sine	\mathbb{R}	$[-1, 1]$
(ii)	cosine	\mathbb{R}	$[-1, 1]$
(iii)	tangent	$\mathbb{R} - \left\{x : x = (2n+1)\frac{\pi}{2}; n \in \mathbb{Z}\right\}$	\mathbb{R}
(iv)	cosecant	$\mathbb{R} - \{x : x = n\pi, n \in \mathbb{Z}\}$	$\mathbb{R} - (-1, 1)$
(v)	secant	$\mathbb{R} - \left\{x : x = (2n+1)\frac{\pi}{2}; n \in \mathbb{Z}\right\}$	$\mathbb{R} - (-1, 1)$
(vi)	cotangent	$\mathbb{R} - \{x : x = n\pi, n \in \mathbb{Z}\}$	\mathbb{R}

1. Inverse function

We know that if function $f: X \rightarrow Y$ such that $y = f(x)$ is **one-one** and **onto**, then we define another function $g: Y \rightarrow X$ such that $x = g(y)$, where $x \in X$ and $y \in Y$, which is also one-one and onto.


In such a case, Domain of $g =$ Range of f

and Range of $g =$ Domain of f
 g is called the inverse of f

$$g = f^{-1}$$

or Inverse of $g = g^{-1} = (f^{-1})^{-1} = f$

The graph of sine function is shown here:



Principal value of branch function \sin^{-1} : It

is a function with domain $[-1, 1]$ and range


$\left[-\frac{3\pi}{2}, -\frac{\pi}{2}\right], \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ or $\left[\frac{\pi}{2}, \frac{3\pi}{2}\right]$ and so on

corresponding to each interval, we get a branch of

the function $\sin^{-1} x$. The branch with range $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$



is called the principal value branch. Thus, $\sin^{-1} : [-1,$

$1] \rightarrow \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.



Principal value branch of function \cos^{-1} : The graph of the function \cos^{-1} is as shown in figure. Domain of the function \cos^{-1} is $[-1, 1]$. Its range in one of the intervals $(-\pi, 0)$, $(0, \pi)$, $(\pi, 2\pi)$, etc. is one-one and onto with the range $[-1, 1]$. The branch with range $(0, \pi)$ is called the principal value branch of the function \cos^{-1} .


Thus, $\cos^{-1} : [-1, 1] \rightarrow [0, \pi]$




Principal value branch function \tan^{-1} : The function \tan^{-1} is defined whose domain is set of real numbers and range is one of the intervals,

$$\left(-\frac{3\pi}{2}, -\frac{\pi}{2}\right), \left(-\frac{\pi}{2}, \frac{\pi}{2}\right), \left(\frac{\pi}{2}, \frac{3\pi}{2}\right), \dots$$

Graph of the function is as shown in the figure:







The branch with range $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ is called the principal value branch of function \tan^{-1} . Thus, $\tan^{-1} : \mathbb{R} \rightarrow \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.

Principal value branch of function cosec⁻¹ : The graph of function cosec⁻¹ is shown in the figure. The cosec⁻¹ is defined on a function whose domain is $\mathbb{R} - (-1, 1)$ and the range is any one of the interval,

$$\left[\frac{-3\pi}{2}, \frac{-\pi}{2}\right] - \{\pi\}, \left[\frac{-\pi}{2}, \frac{\pi}{2}\right] - \{0\}, \left[\frac{\pi}{2}, \frac{3\pi}{2}\right] - \{\pi\}, \dots$$





The function corresponding to the range $\left[\frac{-\pi}{2}, \frac{\pi}{2}\right] - \{0\}$ is called the principal value branch of cosec⁻¹.

$$\text{Thus, } \text{cosec}^{-1} : \mathbb{R} - (-1, 1) \rightarrow \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}.$$


Principal value branch of function sec⁻¹ : The graph of function sec⁻¹ is shown in figure. The sec⁻¹ is defined as a function whose domain $\mathbb{R} - (-1, 1)$ and range is $[-\pi, 0] - \left[-\frac{\pi}{2}, \frac{\pi}{2}\right], [0, \pi] - \left[\frac{\pi}{2}, \frac{3\pi}{2}\right], [\pi, 2\pi] - \left[\frac{3\pi}{2}, \frac{5\pi}{2}\right],$


etc. Function corresponding to range $[0, \pi] - \left[\frac{\pi}{2}, \frac{3\pi}{2}\right]$ is known as the principal value branch of sec⁻¹.

$$\text{Thus, } \text{sec}^{-1} : \mathbb{R} - (-1, 1) \rightarrow [0, \pi] - \left\{\frac{\pi}{2}\right\}$$



The principal value branch of function cot⁻¹ : The graph of function cot⁻¹ is shown below:





The \cot^{-1} function is defined on function whose domain is \mathbb{R} and the range is any of the intervals, $(-\pi, 0)$, $(0, \pi)$, $(\pi, 2\pi)$,

The function corresponding to $(0, \pi)$ is called the principal value branch of the function \cot^{-1} .

Then, $\cot^{-1} : \mathbb{R} \rightarrow (0, \pi)$

The principal value branch of trigonometric inverse functions is as follows:

Inverse Function	Domain	Principal Value Branch
\sin^{-1}	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
\cos^{-1}	$[-1, 1]$	$[0, \pi]$
$\operatorname{cosec}^{-1}$	$\mathbb{R} - (-1, 1)$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$
\sec^{-1}	$\mathbb{R} - (-1, 1)$	$[0, \pi] - \left\{\frac{\pi}{2}\right\}$
\tan^{-1}	\mathbb{R}	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
\cot^{-1}	\mathbb{R}	$(0, \pi)$

Key Facts

- Inverse trigonometric functions are used to find the elevation of sun to the ground. The angle of tilt of the building can be found using inverse trigonometric functions.
- Inverse trigonometric functions help in identifying the angles of bridges to build scale models.

(3) Principal Value:

Numerically smallest angle is known as the principal value.

Finding the principal value: For finding the principal value, following algorithm can be followed :

STEP 1: First draw a trigonometric circle and mark the quadrant in which the angle may lie.

STEP 2: Select anti-clockwise direction for 1st and 2nd quadrants and clockwise direction for 3rd and 4th quadrants.

STEP 3: Find the angles in the first rotation.

STEP 4: Select the numerically least (magnitude wise) angle among these two values. The angle thus found will be the principal value.

STEP 5: In case, two angles one with positive sign and the other with the negative sign qualify for the numerically least angle then, it is the convention to select the angle with positive sign as principal value.

The principal value is never numerically greater than π .

- (4) **To simplify inverse trigonometric expressions, following substitutions can be considered:**

Expression	Substitution
$a^2 + x^2$ or $\sqrt{a^2 + x^2}$	$x = a \tan \theta$ or $x = a \cot \theta$
$a^2 - x^2$ or $\sqrt{a^2 - x^2}$	$x = a \sin \theta$ or $x = a \cos \theta$
$x^2 - a^2$ or $\sqrt{x^2 - a^2}$	$x = a \sec \theta$ or $x = a \operatorname{cosec} \theta$
$\frac{\sqrt{a-x}}{\sqrt{a+x}}$ or $\frac{\sqrt{a+x}}{\sqrt{a-x}}$	$x = a \cos 2\theta$
$\frac{\sqrt{a^2 - x^2}}{\sqrt{a^2 + x^2}}$ or $\frac{\sqrt{a^2 + x^2}}{\sqrt{a^2 - x^2}}$	$x^2 = a^2 \cos 2\theta$

$\sqrt{\frac{x}{a-x}}$ or $\sqrt{\frac{a-x}{x}}$	$x = a \sin^2 \theta$ or $x = a \cos^2 \theta$
$\sqrt{\frac{x}{a+x}}$ or $\sqrt{\frac{a+x}{x}}$	$x = a \tan^2 \theta$ or $x = a \cot^2 \theta$

Note the following and keep them in mind:

- The symbol $\sin^{-1} x$ is used to denote the **smallest angle** whether positive or negative, such that the sine of this angle will give us x . Similarly $\cos^{-1} x$, $\tan^{-1} x$, $\operatorname{cosec}^{-1} x$, $\sec^{-1} x$ and $\cot^{-1} x$ are defined.
- You should note that $\sin^{-1} x$ can be written as **arc sin** x . Similarly, other Inverse Trigonometric Functions can also be written as $\arccos x$, $\arctan x$, $\operatorname{arcsec} x$ etc.
- Keep in mind that these inverse trigonometric relations are **true only in their domains** i.e., they are valid only for some values of ' x ' for which inverse trigonometric functions are well defined.


**Mnemonics**

Inverse trigonometric ratio can be used to find the angle of a right triangle when given two sides of the triangle.

SOH $\theta = \sin^{-1} \frac{\text{opposite}}{\text{hypotenuse}}$

CAH $\theta = \cos^{-1} \frac{\text{adjacent}}{\text{hypotenuse}}$

TOA $\theta = \tan^{-1} \frac{\text{opposite}}{\text{adjacent}}$

**Key Formulæ****TRIGONOMETRIC FORMULAE (ONLY FOR REFERENCE):**➤ **Relation between trigonometric ratios:**

(a) $\tan \theta = \frac{\sin \theta}{\cos \theta}$

(b) $\tan \theta = \frac{1}{\cot \theta}$

(c) $\cot \theta = \frac{\cos \theta}{\sin \theta}$

(d) $\operatorname{cosec} \theta = \frac{1}{\sin \theta}$

(e) $\sec \theta = \frac{1}{\cos \theta}$

➤ **Trigonometric Identities:**

(a) $\sin^2 \theta + \cos^2 \theta = 1$

(b) $\sec^2 \theta = 1 + \tan^2 \theta$

(c) $\operatorname{cosec}^2 \theta = 1 + \cot^2 \theta$

➤ **Addition/subtraction/ formulae & some related results:**

(a) $\sin (A \pm B) = \sin A \cos B \pm \cos A \sin B$

(b) $\cos (A \pm B) = \cos A \cos B \mp \sin A \sin B$

(c) $\cos (A + B) \cos (A - B) = \cos^2 A - \sin^2 B = \cos^2 B - \sin^2 A$

(d) $\sin (A + B) \sin (A - B) = \sin^2 A - \sin^2 B = \cos^2 B - \cos^2 A$

(e) $\tan (A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$

(f) $\cot (A \pm B) = \frac{\cot B \cot A \mp 1}{\cot B \pm \cot A}$

➤ **Multiple angle formulae involving A , $2A$ & $3A$:**

(a) $\sin 2A = 2 \sin A \cos A$

(b) $\sin A = 2 \sin \frac{A}{2} \cos \frac{A}{2}$

(c) $\cos 2A = \cos^2 A - \sin^2 A$

(d) $\cos A = \cos^2 \frac{A}{2} - \sin^2 \frac{A}{2}$

(e) $\cos 2A = 2 \cos^2 A - 1$

(f) $2 \cos^2 A = 1 + \cos 2A$

(g) $\cos 2A = 1 - 2 \sin^2 A$

(h) $2 \sin^2 A = 1 - \cos 2A$

(i) $\sin 2A = \frac{2 \tan A}{1 + \tan^2 A}$

(j) $\cos 2A = \frac{1 - \tan^2 A}{1 + \tan^2 A}$

(k) $\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$

(l) $\sin 3A = 3 \sin A - 4 \sin^3 A$

(m) $\cos 3A = 4 \cos^3 A - 3 \cos A$

(n) $\tan 3A = \frac{3 \tan A - \tan^3 A}{1 - 3 \tan^2 A}$

➤ Transformation of sums/differences into products & vice-versa:

(a) $\sin C + \sin D = 2 \sin \frac{C+D}{2} \cos \frac{C-D}{2}$

(b) $\sin C - \sin D = 2 \cos \frac{C+D}{2} \sin \frac{C-D}{2}$

(c) $\cos C + \cos D = 2 \cos \frac{C+D}{2} \cos \frac{C-D}{2}$

(d) $\cos C - \cos D = -2 \sin \frac{C+D}{2} \sin \frac{C-D}{2}$

(e) $2 \sin A \cos B = \sin(A+B) + \sin(A-B)$

(f) $2 \cos A \sin B = \sin(A+B) - \sin(A-B)$

(g) $2 \cos A \cos B = \cos(A+B) + \cos(A-B)$

(h) $2 \sin A \sin B = \cos(A-B) - \cos(A+B)$

➤ Relations in different measures of Angle:

(a) Angle in Radian Measure = (Angle in degree measure) $\times \frac{\pi}{180^\circ}$ rad

(b) Angle in Degree Measure = (Angle in radian measure) $\times \frac{180^\circ}{\pi}$

(c) θ (in radian measure) = $\frac{l}{r} = \frac{\text{arc}}{\text{radius}}$

Also following are of importance as well:

(a) 1 right angle = 90°

(b) $1^\circ = 60', 1' = 60''$

(c) $1^\circ = \frac{\pi}{180^\circ} = 0.01745$ radians (Approx.)

(d) 1 radian = $57^\circ 17' 45''$ or 206265 seconds.

➤ General Solutions:

(a) $\sin x = \sin y$ or, $x = n\pi + (-1)^n y$, where $n \in \mathbb{Z}$.

(b) $\cos x = \cos y$ or, $x = 2n\pi \pm y$, where $n \in \mathbb{Z}$.

(c) $\tan x = \tan y$ or, $x = n\pi + y$, where $n \in \mathbb{Z}$.

➤ Relation in Degree & Radian Measures:

Angles in Degree	0°	30°	45°	60°	90°	180°	270°	360°
Angles in Radian	0°	$\left(\frac{\pi}{6}\right)$	$\left(\frac{\pi}{4}\right)$	$\left(\frac{\pi}{3}\right)$	$\left(\frac{\pi}{2}\right)$	(π)	$\left(\frac{3\pi}{2}\right)$	(2π)

➤ Trigonometric Ratio of Standard Angles:

Degree	0°	30°	45°	60°	90°
$\sin x$	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1
$\cos x$	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0
$\tan x$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	∞
$\cot x$	∞	$\sqrt{3}$	1	$\frac{1}{\sqrt{3}}$	0
$\operatorname{cosec} x$	∞	2	$\sqrt{2}$	$\frac{2}{\sqrt{3}}$	1
$\sec x$	1	$\frac{2}{\sqrt{3}}$	$\sqrt{2}$	2	∞

Trigonometric Ratios of Allied Angles:

Angles (\rightarrow)	$\frac{\pi}{2} - \theta$	$\frac{\pi}{2} + \theta$	$\pi - \theta$	$\pi + \theta$	$\frac{3\pi}{2} - \theta$	$\frac{3\pi}{2} + \theta$	$2\pi - \theta$ or $-\theta$	$2\pi + \theta$
T - Ratios (\downarrow)								
sin	cos θ	cos θ	sin θ	- sin θ	- cos θ	- cos θ	- sin θ	sin θ
cos	sin θ	- sin θ	- cos θ	- cos θ	- sin θ	sin θ	cos θ	cos θ
tan	cot θ	- cot θ	- tan θ	tan θ	cot θ	- cot θ	- tan θ	tan θ
cot	tan θ	- tan θ	- cot θ	cot θ	tan θ	- tan θ	- cot θ	cot θ
sec	cosec θ	- cosec θ	- sec θ	- sec θ	- cosec θ	cosec θ	sec θ	sec θ
cosec	sec θ	sec θ	cosec θ	- cosec θ	- sec θ	- sec θ	- cosec θ	cosec θ

UNIT - II : ALGEBRA

CHAPTER-3

MATRICES

Topic-1**Matrices and Operations**

Concepts Covered • Basic concept of matrices,
• Types of matrices, • Operations on matrices

**Revision Notes****1. MATRIX - BASIC INTRODUCTION:**

A matrix is an ordered rectangular **array** of numbers (real or complex) or functions which are known as elements or the entries of the matrix. It is denoted by the uppercase letters i.e. A, B, C etc.

Key Words

Array: An array is a rectangular arrangement of objects in form of rows (horizontal) and columns (vertical). Everyday example of arrays include a muffin tray and an egg tray.

Consider a matrix A given as,

Here in matrix A the horizontal lines of elements are said to constitute **rows** and vertical lines of elements are said to constitute **columns** of the matrix. Thus, matrix A has m **rows** and n **columns**. The array is enclosed by square brackets [], the parentheses () or the double vertical bars || ||.

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1j} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2j} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots & & \vdots \\ a_{i1} & a_{i2} & \dots & a_{ij} & \dots & a_{in} \\ \vdots & \vdots & & \vdots & & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mj} & \dots & a_{mn} \end{bmatrix}_{m \times n}$$

- A matrix having m rows and n columns is called a matrix of order $m \times n$ (read as 'm by n' matrix). A matrix A of order $m \times n$ is depicted as $A = [a_{ij}]_{m \times n}$; $i, j \in N$.
- Also in general, a_{ij} means an element lying in the i^{th} row and j^{th} column.
- Number of elements in the matrix $A = [a_{ij}]_{m \times n}$ is given as mn .

2. TYPES OF MATRICES:

- (i) **Column matrix:** A matrix having only one column is called a **column matrix** or **column vector**.

$$\text{e.g.: } \begin{bmatrix} 0 \\ 1 \\ -2 \end{bmatrix}_{3 \times 1}, \begin{bmatrix} 4 \\ 5 \end{bmatrix}_{2 \times 1}$$

General notation : $A = [a_{ij}]_{m \times n}$



Key Fact

- The term matrix was introduced by the 19th century English Mathematician James Sylvester, but it was his friend the Mathematician Arthur Cayley who developed the algebraic aspect of matrices in two papers in the 1850s.

(ii) **Row matrix:** A matrix having only one row is called a **row matrix** or **row vector**.

General notation : $A = [a_{ij}]_{1 \times n}$

(iii) **Square matrix:** It is a matrix in which the number of rows is equal to the number of columns *i.e.*, an $n \times n$ matrix is said to constitute a square matrix of order $n \times n$ and is known as a **square matrix of order 'n'**.

General notation : $A = [a_{ij}]_{n \times n}$

(iv) **Diagonal matrix:** A square matrix $A = [a_{ij}]_{m \times m}$ is said to be a **diagonal matrix** if all the elements, except those in the leading diagonal are zero *i.e.*, $a_{ij} = 0$, for all $i \neq j$.

(v) **Scalar matrix:** A diagonal matrix $A = [a_{ij}]_{m \times m}$ is said to be a **scalar matrix** if its diagonal elements are equal. *i.e.*,

$$a_{ij} = \begin{cases} 0, & \text{when } i \neq j \\ k, & \text{when } i = j \text{ for some constant } k \end{cases}$$

(vi) **Unit or Identity matrix:** A square matrix $A = [a_{ij}]_{m \times m}$ is said to be an **identity matrix** if $a_{ij} = \begin{cases} 1, & \text{if } i = j \\ 0, & \text{if } i \neq j \end{cases}$

A **unit matrix** can also be defined as the **scalar matrix** in which all diagonal elements are equal to **unity**. We denote the identity matrix of order m by I_m or I .

(vii) **Zero matrix or Null matrix:** A matrix is said to be a **zero matrix** or **null matrix** if each of its elements is '0'.

(viii) **Horizontal matrix:** A $m \times n$ matrix is said to be a **horizontal matrix** if $m < n$.

(ix) **Vertical matrix:** A $m \times n$ matrix is said to be a **vertical matrix** if $m > n$.

3. EQUALITY OF MATRICES:

Two matrices A and B are said to be equal and written as $A = B$, if they are of the **same order** and their **corresponding elements are identical** *i.e.* $a_{ij} = b_{ij}$ *i.e.*, $a_{11} = b_{11}$, $a_{22} = b_{22}$, $a_{32} = b_{32}$ etc.

4. ADDITION OF MATRICES:

If A and B are two $m \times n$ matrices, then another $m \times n$ matrix obtained by adding the corresponding elements of the matrices A and B is called the sum of the matrices A and B and is denoted by ' $A + B$ '.

Thus if $A = [a_{ij}]$, $B = [b_{ij}]$, or $A + B = [a_{ij} + b_{ij}]$.

5. MULTIPLICATION OF A MATRIX BY A SCALAR:

If a $m \times n$ matrix A is multiplied by a scalar k (say), then the new kA matrix is obtained by multiplying each element of matrix A by scalar k . Thus, if $A = [a_{ij}]$ and it is multiplied by a scalar k , then $kA = [ka_{ij}]$, *i.e.* $A = [a_{ij}]$ or $kA = [ka_{ij}]$.

6. MULTIPLICATION OF TWO MATRICES:

Let $A = [a_{ij}]$ be a $m \times n$ matrix and $B = [b_{jk}]$ be a $n \times p$ matrix such that the number of columns in A is equal to the number of rows in B , then the $m \times p$ matrix $C = [c_{ik}]$ such that $[c_{ik}] = \sum_{j=1}^n a_{ij} b_{jk}$ is said to be the product of the matrices A and B in that order and it is denoted by AB *i.e.* " $C = AB$ ".

Properties of matrix multiplication:

- Note that the product AB is defined only when the number of columns in matrix A is equal to the number of rows in matrix B .
- If A and B are $m \times n$ and $n \times p$ matrices, respectively, then the matrix AB will be an $m \times p$ matrix *i.e.*, order of matrix AB will be $m \times p$.
- If A is a $m \times n$ matrix and both AB as well as BA are defined, then B will be a $n \times m$ matrix.
- If A is a $n \times n$ matrix and I_n be the unit matrix of order n , then $A I_n = I_n A = A$.
- Matrix multiplication is not commutative.

7. IDEMPOTENT MATRIX:

A square matrix A is said to be an idempotent matrix if $A^2 = A$.

For example,

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad A = \begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{bmatrix}$$

are idempotent matrices.

8. TRANSPOSE OF A MATRIX:

If $A = [a_{ij}]_{m \times n}$ be a $m \times n$ matrix, then the matrix obtained by interchanging the rows and columns of matrix A is said to be a **transpose of matrix** A . The transpose of A is denoted by A' or A^T *i.e.*, if $A^T = [a_{ji}]_{n \times m}$.

PROPERTIES OF TRANSPOSE OF MATRICES:

- $(A + B)^T = A^T + B^T$
- $(A^T)^T = A$
- $(kA)^T = kA^T$, where k is any constant
- $(AB)^T = B^T A^T$
- $(ABC)^T = C^T B^T A^T$



Mnemonics

Types of Matrices

Ram	Charan	Says	Drink	Sprite
↓	↓	↓	↓	↓
Row Matrix	Column Matrix	Square Matrix	Diagonal Matrix	Scalar Matrix
and	Nescafe	Ice Tea		
	↓	↓	↓	
	Null Matrix	Identity Matrix	Triangular Matrix	

Matrix Multiplication

No. of columns of first matrix = No. of rows of second matrix

$$\begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix} \begin{bmatrix} u & v \\ w & x \\ y & z \end{bmatrix}$$

$$R \times C \quad R \times C$$

Class Representative

Topic-2

Symmetric, Skew Symmetric and Invertible Matrices

Concepts Covered • Symmetric Matrix, • Skew Symmetric Matrix, • Invertible Matrix
• Uniqueness Theorem



Revision Notes

Symmetric matrix: A square matrix $A = [a_{ij}]$ is said to be a **symmetric matrix** if $A^T = A$. i.e., if $A = [a_{ij}]$, then $A^T = [a_{ji}] = [a_{ij}]$ or $A^T = A$.

For example:

$$\begin{bmatrix} a & h & g \\ h & b & f \\ g & f & c \end{bmatrix}, \begin{bmatrix} 2+i & 1 & 3 \\ 1 & 2 & 3+2i \\ 3 & 3+2i & 4 \end{bmatrix}$$

Skew symmetric matrix: A square matrix $A = [a_{ij}]$ is said to be a **skew symmetric matrix** if

$A^T = -[A]$ i.e., if $A = [a_{ij}]$, then $A^T = [a_{ji}] = -[a_{ij}]$ or $A^T = -A$.

For example: $\begin{bmatrix} 0 & 1 & -5 \\ -1 & 0 & 5 \\ 5 & -5 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}$

Orthogonal matrix: A matrix A is said to be **orthogonal** if $A.A^T = I$, where A^T is transpose of A .

Invertible Matrix: An invertible matrix is a matrix for which matrix inversion operation exists, given that it satisfies the requisite conditions. Any given square matrix A of order $n \times n$ is called invertible if there exists another $n \times n$ square matrix B such that, $AB = BA = I_n$, where I_n is on identity matrix of order $n \times n$.

Example: Let matrix $A = \begin{bmatrix} 1 & 2 \\ 2 & 5 \end{bmatrix}$ and matrix $B =$

$$\begin{bmatrix} 5 & -2 \\ -2 & 1 \end{bmatrix}$$

Now, $AB = \begin{bmatrix} 1 & 2 \\ 2 & 5 \end{bmatrix} \begin{bmatrix} 5 & -2 \\ -2 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

and $BA = \begin{bmatrix} 5 & -2 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 2 & 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

Hence, $A^{-1} = B$ and B is called the inverse of A .

So, A can also be the inverse of B or $B^{-1} = A$.

Uniqueness of Inverse of Matrix

If there exists an inverse of a square matrix, it is always unique.

Proof: Let A be a square matrix of order $n \times n$. Let us assume matrices B and C be inverses of matrix A .

Now, $AB = BA = I$, since B is the inverse of matrix A .

Similarly, $AC = CA = I$

But, $B = BI = B(AC) = (BA)C = IC = C$

This proves $B = C$, or B and C are the same matrices.

Key Facts

- Note that $[a_{ji}] = -[a_{ij}]$ or $[a_{ii}] = -[a_{ii}]$ or $2[a_{ii}] = 0$ (Replacing j by i). i.e., all the diagonal elements in a skew symmetric matrix are zero.
- For any matrices, AA^T and $A^T A$ are symmetric matrices.
- For a square matrix A , the matrix $A + A^T$ is a symmetric matrix and $A - A^T$ is always a skew-symmetric matrix.
- Also note that any square matrix can be expressed as the sum of a symmetric and a skew symmetric matrix i.e., $A = P + Q$ where $P = \frac{A + A^T}{2}$ is a symmetric matrix and $Q = \frac{A - A^T}{2}$ is a skew symmetric matrix.

CHAPTER-4

DETERMINANTS

Topic-1

Determinants, Minors & Co-factors

- Concepts Covered**
- Determinant value of a matrix,
 - Co-factor and Minor of a matrix
 - Inverse of matrix using Adjoint method, • Area of triangle with the help of determinant



Revision Notes

Determinants, Minors & Co-factors

- (a) **Determinant:** A unique number (real or complex) can be associated to every square matrix $A = [a_{ij}]$ of order m . This number is called the determinant of the square matrix A , where $a_{ij} = (i, j)^{\text{th}}$ element of A .

For instance, if $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ then, determinant

of matrix A is written as $|A| = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = \det(A)$

and its value is given by $ad - bc$.

- (b) **Minors:** Minors of an element a_{ij} of a determinant (or a determinant corresponding to matrix A) is the determinant obtained by deleting its i^{th} row and j^{th} column in which a_{ij} lies. Minor of a_{ij} is denoted by M_{ij} . Hence, we can get 9 minors corresponding to the 9 elements of a third order (i.e., 3×3) determinant.
- (c) **Co-factors:** Cofactor of an element a_{ij} denoted by A_{ij} is defined by $A_{ij} = (-1)^{(i+j)} M_{ij}$, where M_{ij} is minor of a_{ij} . Sometimes C_{ij} is used in place of A_{ij} to denote the co-factor of element a_{ij} .

1. ADJOINT OF A SQUARE MATRIX:

Let $A = [a_{ij}]$ be a square matrix. Also, assume $B = [A_{ij}]$, where A_{ij} is the cofactor of the elements a_{ij} in matrix A . Then the transpose B^T of matrix B is called the **adjoint of matrix A** and it is denoted by " $\text{adj}(A)$ ".

To find adjoint of a 2×2 matrix: Follow this,

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \text{ or } \text{adj } A = \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

For example, consider a square matrix of order 3 as

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 2 & 0 & 5 \end{bmatrix}, \text{ then in order to find the adjoint}$$

matrix A , we find a matrix B (formed by the co-factors of elements of matrix A as mentioned above in the definition)

$$\text{i.e., } B = \begin{bmatrix} 15 & -2 & -6 \\ -10 & -1 & 4 \\ -1 & 2 & -1 \end{bmatrix}.$$

$$\text{Hence, } \text{adj } A = B^T = \begin{bmatrix} 15 & -10 & -1 \\ -2 & -1 & 2 \\ -6 & 4 & -1 \end{bmatrix}$$

2. SINGULAR MATRIX AND NON-SINGULAR MATRIX:

(a) **Singular matrix:** A square matrix A is said to be singular if $|A| = 0$ i.e., its determinant is zero.

$$\text{e.g. } A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 12 \\ 1 & 1 & 3 \end{bmatrix}$$

$$= 1(15 - 12) - 2(12 - 12) + 3(4 - 5) \\ = 3 - 0 - 3 = 0$$

$\therefore A$ is singular matrix.

(b) **Non-singular matrix:** A square matrix A is said to be non-singular if $|A| \neq 0$.

$$\text{e.g. } A = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

$$= 0(0 - 1) - 1(0 - 1) + 1(1 - 0)$$

$$= 0 + 1 + 1 = 2 \neq 0$$

$\therefore A$ is non-singular matrix.

- A square matrix A is **invertible** if and only if A is **non-singular**.

3. ALGORITHM TO FIND A^{-1} BY

DETERMINANT METHOD:

STEP 1: Find $|A|$.

STEP 2: If $|A| = 0$, then, write “ A is a singular matrix and hence not invertible”. Else write “ A is a non-singular matrix and hence invertible”.

STEP 3: Calculate the co-factors of elements of matrix A .

STEP 4: Write the matrix of co-factors of elements of A and then obtain its transpose to get $\text{adj.}A$ (i.e., adjoint A).

STEP 5: Find the inverse of A by using the relation

$$A^{-1} = \frac{1}{|A|}(\text{adj.}A).$$

4. AREA OF TRIANGLE:

Area of a triangle whose vertices are (x_1, y_1) , (x_2, y_2) and (x_3, y_3) is given by,

$$\Delta = \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} \text{ sq. units}$$

- Since area is a positive quantity, we take absolute value of the determinant.
- If the points (x_1, y_1) , (x_2, y_2) and (x_3, y_3) are collinear, then $\Delta = 0$.
- The equation of a line passing through the points (x_1, y_1) and (x_2, y_2) can be obtained by the expression given here:

$$\begin{vmatrix} x & y & 1 \\ x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \end{vmatrix} = 0$$



Key Fact

- In mathematics, the determinant is a scalar value that is a function of the entries of a square matrix.

Topic-2

Solutions of System of Linear Equations

Concepts Covered • Unique Solution, • Consistent System, • Inconsistent System



Revision Notes

SOLVING SYSTEM OF EQUATIONS BY MATRIX METHOD [INVERSE MATRIX METHOD]

(a) **Homogeneous and Non-homogeneous system:** A system of equations $AX = B$ is said to be a homogeneous system if $B = O$. Otherwise it is called a non-homogeneous system of equations.

$$a_1x + b_1y + c_1z = d_1,$$

$$a_2x + b_2y + c_2z = d_2,$$

$$a_3x + b_3y + c_3z = d_3$$

STEP 1: Assume

$$A = \begin{bmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{bmatrix}, B = \begin{bmatrix} d_1 \\ d_2 \\ d_3 \end{bmatrix} \text{ and } X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}.$$

STEP 2: Find $|A|$. Now there may be following situations :

(i) $|A| \neq 0 \Rightarrow A^{-1}$ exists. It implies that the given system of equations is consistent and therefore, the system has **unique solution**. In that case, write

$$\Rightarrow \begin{matrix} AX = B \\ X = A^{-1}B \end{matrix}$$

$$\left[\text{where } A^{-1} = \frac{1}{|A|}(\text{adj.}A) \right]$$

Then by using the definition of equality of matrices, we can get the values of x , y and z .

- (ii) $|A| = 0 \Rightarrow A^{-1}$ does not exist. It implies that the given system of equations may be **consistent** or **inconsistent**.



Key Words

Consistent System: A system is considered to be consistent if it has at least one solution.

Inconsistent System: If a system has no solution, it is said to be inconsistent.

In order to check proceed as follow:

\Rightarrow Find $(adj A) B$. Now, we may have either $(adj A) B \neq O$ or $(adj A) B = O$.

- If $(adj A)B = O$, then the given system may be consistent or inconsistent.

To check, put $z = k$ in the given equations and proceed in the same manner in the new two variables system of equations assuming $d_i - c_i k$, $1 \leq i \leq 3$ as constant.

- And if $(adj A) B \neq O$, then the given system is inconsistent with no solutions.



Mnemonics-1

Inverse of a Square Matrix

Det mins tra

↓ ↓ ↓
 Determinant Minors along with signs Transpose



Mnemonics-2

Singular Matrix

A square matrix is said to be singular matrix if determinant of matrix denoted by $|A|$ is zero otherwise it is non singular matrix

Inverse Of a Matrix

Determinant

"a **D**etermined **A**rtist Can become a **S**inger,
 if he is **O**ptimistic. (Zero)
 "a **D**etermined **A**rtist Can **N**ever be **S**inger
 if he is **N**ot **O**ptimistic
 Non Singular
 if $|A| = 0$, then A is Singular Otherwise,
 A is non-Singular

$\neq 0$ (Zero)

"If **D**etermined **A**rtist is **N**ot **O**ptimistic

then **A**DJust **B**y **I**nverse

Adjoint By Inverse

"Musicians"
 ↓
 Matrix

Determinant

"A is (non-singular) i.e. $|A| \neq 0$ then

$$A^{-1} = \frac{1}{|A|} \cdot (adj A)$$

Interpretation:

Singular & Non Singular Matrix -

if $|A| = 0$, then A is singular. Otherwise A is non-singular

Inverse of a Matrix-

Inverse of a Matrix exists if A is non-singular i.e. $|A| \neq 0$, and is given by

$$A^{-1} = \frac{1}{|A|} adj A$$

UNIT – III: CALCULUS**CHAPTER-5****CONTINUITY & DIFFERENTIABILITY****Topic-1****Continuity**

Concepts Covered • Left hand Limit,
• Right Hand Limit

**Revision Notes****FORMULAE FOR LIMITS:**

(a) $\lim_{x \rightarrow 0} \cos x = 1$

(b) $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$

(c) $\lim_{x \rightarrow 0} \frac{\tan x}{x} = 1$

(d) $\lim_{x \rightarrow 0} \frac{\sin^{-1} x}{x} = 1$

(e) $\lim_{x \rightarrow 0} \frac{\tan^{-1} x}{x} = 1$ (f) $\lim_{x \rightarrow 0} \frac{a^x - 1}{x} = \log_e a, a > 0$

(g) $\lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1$ (h) $\lim_{x \rightarrow 0} \frac{\log_e(1+x)}{x} = 1$

(i) $\lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = na^{n-1}$

⇒ For a function $f(x)$, $\lim_{x \rightarrow m} f(x)$ exists if

$$\lim_{x \rightarrow m^-} f(x) = \lim_{x \rightarrow m^+} f(x).$$

⇒ A function $f(x)$ is continuous at a point $x = m$ if, $\lim_{x \rightarrow m^-} f(x) = \lim_{x \rightarrow m^+} f(x) = f(m)$, where

$\lim_{x \rightarrow m^-} f(x)$ is **Left Hand Limit** of $f(x)$ at $x = m$

and $\lim_{x \rightarrow m^+} f(x)$ is **Right Hand Limit** of $f(x)$ at $x = m$. Also $f(m)$ is the value of function $f(x)$ at $x = m$.

⇒ A function $f(x)$ is continuous at $x = m$ (say) if, $f(m) = \lim_{x \rightarrow m} f(x)$ i.e., a function is continuous at a point in its domain if the **limit value of the function** at that point **equals** the value of the function at the same point.

⇒ For a continuous function $f(x)$ at $x = m$, $\lim_{x \rightarrow m} f(x)$ can be directly obtained by evaluating $f(m)$.

⇒ Indeterminate forms or meaningless forms:

$$\frac{0}{0}, \frac{\infty}{\infty}, 0 \times \infty, \infty - \infty, 1^\infty, 0^0, \infty^0.$$

Topic-2**Differentiability**

Concepts Covered • Left Hand Derivative, • Right Hand Derivative,
• Relation between Continuity and Differentiability

Derivative of Some Standard Functions:

(a) $\frac{d}{dx}(x^n) = nx^{n-1}$

(b) $\frac{d}{dx}(k) = 0$, where k is any constant

(c) $\frac{d}{dx}(a^x) = a^x \log_e a, a > 0$

(d) $\frac{d}{dx}(e^x) = e^x$

(e) $\frac{d}{dx}(\log_a x) = \frac{1}{x \log_e a} = \frac{1}{x} \log_a e$

(f) $\frac{d}{dx}(\log_e x) = \frac{1}{x}$

(g) $\frac{d}{dx}(\sin x) = \cos x$

(h) $\frac{d}{dx}(\cos x) = -\sin x$

(i) $\frac{d}{dx}(\tan x) = \sec^2 x$

(j) $\frac{d}{dx}(\sec x) = \sec x \tan x$

(k) $\frac{d}{dx}(\cot x) = -\operatorname{cosec}^2 x$

(l) $\frac{d}{dx}(\operatorname{cosec} x) = -\operatorname{cosec} x \cot x$

(m) $\frac{d}{dx}(\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}, x \in (-1, 1)$

(n) $\frac{d}{dx}(\cos^{-1} x) = -\frac{1}{\sqrt{1-x^2}}, x \in (-1, 1)$

(o) $\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}, x \in \mathbb{R}$

(p) $\frac{d}{dx}(\cot^{-1} x) = -\frac{1}{1+x^2}, x \in \mathbb{R}$

(q) $\frac{d}{dx}(\sec^{-1} x) = \frac{1}{x\sqrt{x^2-1}}, \text{ where } x \in (-\infty, -1) \cup (1, \infty)$

(r) $\frac{d}{dx}(\operatorname{cosec}^{-1} x) = -\frac{1}{x\sqrt{x^2-1}}, \text{ where } x \in (-\infty, -1) \cup (1, \infty)$

Following derivatives should also be memorized by you for quick use:

(i) $\frac{d}{dx}(\sqrt{x}) = \frac{1}{2\sqrt{x}}$

(ii) $\frac{d}{dx}\left(\frac{1}{x}\right) = -\frac{1}{x^2}$

➤ **Left Hand Derivative of $f(x)$ at $x = m$,**

$$Lf'(m) = \lim_{x \rightarrow m^-} \frac{f(x) - f(m)}{x - m} \text{ and,}$$

Right Hand Derivative of $f(x)$ at $x = m$,

$$Rf'(m) = \lim_{x \rightarrow m^+} \frac{f(x) - f(m)}{x - m}$$

For a function to be differentiable at a point, LHD and RHD at that point should be equal.

➤ **Derivative of y w.r.t. x :** $\frac{dy}{dx} = \lim_{\delta x \rightarrow 0} \frac{\delta y}{\delta x}$.

Also, for very-very small value h ,

$$f'(x) = \frac{f(x+h) - f(x)}{h}, \text{ (as } h \rightarrow 0)$$

Relation between Continuity and Differentiability:

- (i) If a function is differentiable at a point, it is continuous at that point as well.
- (ii) If a function is not differentiable at a point, it may or may not be continuous at that point.
- (iii) If a function is continuous at a point, it may or may not be differentiable at that point.
- (iv) If a function is **discontinuous** at a point, it is not be differentiable at that point.

Rules of Derivatives:

➤ Product or Leibniz's rule of derivatives:

$$\frac{d}{dx}(uv) = u \frac{d}{dx}(v) + v \frac{d}{dx}(u)$$

➤ Quotient Rule of derivatives:

$$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{d}{dx}(u) - u \frac{d}{dx}(v)}{v^2} = \frac{vu' - uv'}{v^2}$$



Key Word

Discontinuous Function: A discontinuous function is a function in algebra that has a point where either the function is not defined at that point or the LHL and RHL of the function are equal but not equal to the value of the function at that point or the limit of the function does not exist at the given point.



Key Facts

- $f(x) = 0$ is a continuous function because it is an unbroken line, without holes or jumps.
- If $f(0) = \infty$, then function is continuous at 0.
- All polynomial functions are continuous functions.



Mnemonics

Quotient Rule of Derivative

Ho D Hi Minus Hi D Ho Over Ho Ho

In mathematical notation, $\frac{\text{Ho D Hi} - \text{Hi D Ho}}{\text{ho ho}}$

where, Ho \rightarrow function in numerator

Hi \rightarrow function in denominator

D \rightarrow derivative of

CHAPTER-6

APPLICATIONS OF DERIVATIVES

Topic-1 Rate of Change of Bodies



Revision Notes

Interpretation of $\frac{dy}{dx}$ as a rate measure:

If two variables x and y are varying with respect to another variables say t , i.e., if $x = f(t)$, then by the Chain Rule, we have

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt}, \frac{dx}{dt} \neq 0$$

Thus, the rate of change of y with respect to x can be calculated by using the rate of change of y and that of x both with respect to t .

Also, if y is a function of x and they are related as $y = f(x)$ then, $f'(x)$, i.e., represents the rate of change of y with respect to x at the instant when $x = \alpha$.

Topic-2

Increasing/Decreasing Functions

Concepts Covered • Increasing function, • Decreasing function, • Constant function
• Monotonic function



Revision Notes

1. A function $f(x)$ is said to be an increasing function in $[a, b]$, if as x increases, $f(x)$ also increases i.e., if $\alpha, \beta \in [a, b]$ and $\alpha > \beta$, $f(\alpha) > f(\beta)$.

If $f'(x) \geq 0$ lies in (a, b) , then $f(x)$ is an increasing function in $[a, b]$, provided $f(x)$ is continuous at $x = a$ and $x = b$.

2. A function $f(x)$ is said to be a **decreasing function** in $[a, b]$, if, as x increases, $f(x)$ decreases i.e., if $\alpha, \beta \in [a, b]$ and $\alpha > \beta \Rightarrow f(\alpha) < f(\beta)$.

If $f'(x) \leq 0$ lies in (a, b) , then $f(x)$ is a decreasing function in $[a, b]$ provided $f(x)$ is continuous at $x = a$ and $x = b$.

⇒ A function $f(x)$ is a **constant function** in $[a, b]$ if $f'(x) = 0$ for each $x \in (a, b)$.

⇒ By **monotonic function** $f(x)$ in interval I , we mean that f is either **only increasing** in I or **only decreasing** in I .

3. **Finding the intervals of increasing and/or decreasing of a function:**

ALGORITHM

STEP 1: Consider the function $y = f(x)$.

STEP 2: Find $f'(x)$.

STEP 3: Put $f'(x) = 0$ and solve to get the critical point(s).

STEP 4: The value(s) of x for which $f'(x) > 0$, $f(x)$ is increasing; and the value(s) of x for which $f'(x) < 0$, $f(x)$ is decreasing.

Topic-3**Maxima and Minima**

Concepts Covered • Local Maxima, • Local Minima, • Absolute Maxima, • Absolute Minima, • First derivative test, • Second derivative test

**Revision Notes****1. Understanding maxima and minima:**

Consider $y = f(x)$ be a well defined function on an interval I , then

**Key Word**

Interval: In mathematics, an interval is a set of real numbers between two given numbers called end points of the interval.

- (a) f is said to have a **maximum value** in I , if there exists a point c in I such that $f(c) > f(x)$, for all $x \in I$. The value corresponding to $f(c)$ is called the maximum value of f in I and the point c is called the **point of maximum value of f in I** .
- (b) f is said to have a minimum value in I , if there exists a point c in I such that $f(c) < f(x)$, for all $x \in I$. The value corresponding to $f(c)$ is called the minimum value of f in I and the point c is called the **point of minimum value of f in I** .
- (c) f is said to have an **extreme value** in I , if there exists a point c in I such that $f(c)$ is either a maximum value or a minimum value of f in I . The value $f(c)$ in this case, is called an extreme value of f in I and the point c called an **extreme point**.

2. Let f be a real valued function and also take a point c from its domain, then

- (i) c is called a **point of local maxima** if there exists a number $h > 0$ such that $f(c) > f(x)$, for all x in $(c - h, c + h)$. The value $f(c)$ is called the **local maximum value of f** .
- (ii) c is called a **point of local minima** if there exists a number $h > 0$ such that $f(c) < f(x)$, for all x in $(c - h, c + h)$. The value $f(c)$ is called the **local minimum value of f** .

3. Critical points

It is a point c (say) in the domain of a function $f(x)$ at which either $f'(x)$ vanishes i.e., $f'(c) = 0$ or f is not differentiable.

4. First Derivative Test:

Consider $y = f(x)$ be a well defined function on an open interval I . Now proceed as have been mentioned in the following algorithm:

STEP 1: Find $\frac{dy}{dx}$.

STEP 2: Find the critical point(s) by putting $\frac{dy}{dx} = 0$.

Suppose $c \in I$ (where I is the interval) be any critical point

and f be continuous at this point c . Then we may have following situations:

- ⇒ $\frac{dy}{dx}$ changes sign from **positive to negative** as x increases through c , then the function attains a **local maximum** at $x = c$.
- ⇒ $\frac{dy}{dx}$ changes sign from **negative to positive** as x increases through c , then the function attains a **local minimum** at $x = c$.
- ⇒ $\frac{dy}{dx}$ **does not change sign** as x increases through c , then $x = c$ is **neither** a point of **local maximum nor** a point of **local minimum**. Rather in this case, the point $x = c$ is called the **point of inflection**.

5. Second Derivative Test:

Consider $y = f(x)$ be a well defined function on an open interval I and twice differentiable at a point c in the interval. Then we observe that:

- ⇒ $x = c$ is a point of local maxima if $f'(c) = 0$ and $f''(c) < 0$.
The value $f(c)$ is called the local maximum value of f .
- ⇒ $x = c$ is a point of local minima if $f'(c) = 0$ and $f''(c) > 0$.
The value $f(c)$ is called the local minimum value of f .

This test fails if $f'(c) = 0$ and $f''(c) = 0$. In such a case, we use **first derivative test** as discussed above.

6. Absolute maxima and absolute minima:

If f is a continuous function on a **closed interval I** , then f has the absolute maximum value and f attains it atleast once in I . Also f has the absolute minimum value and the function attains it atleast once in I .

ALGORITHM

STEP 1: Find all the critical points of f in the given interval, i.e., find all the points x where either $f'(x) = 0$ or f is not differentiable.

STEP 2: Take the end points of the given interval.

STEP 3: At all these points (i.e., the points found in STEP 1 and STEP 2) calculate the values of f .

STEP 4: Identify the maximum and minimum value of f out of the values calculated in STEP 3. This maximum value will be the **absolute maximum value** of f and the minimum value will be the **absolute minimum value** of the function f .

Absolute maximum value is also called as **global maximum value** or **greatest value**. Similarly absolute minimum value is called as **global minimum value** or the **least value**.

CHAPTER-7

INTEGRALS

Topic-1

Indefinite Integral

Concepts Covered • Meaning of Integral of function • Integration by Substitution
• Integration by partial fraction • Integration by parts
• Formulae for Indefinite Integral



Revision Notes

➤ Meaning of Integral of Function

If differentiation of a function $F(x)$ is $f(x)$ i.e., if $\frac{d}{dx}$

$[F(x)] = f(x)$, then we say that one integral or primitive or anti-derivative of $f(x)$ is $F(x)$ and in symbols, we write,

$$\int f(x)dx = F(x) + C.$$

Therefore, we can say that integration is the inverse process of differentiation.

➤ Methods of Integration

(a) Integration by Substitution Method :

In this method, we change the integral $\int f(x)dx$

where independent variable is x , to another integral in which independent variable is t (say) different from x such that x and t are related by $x = g(t)$.

$$\text{i.e., } \int f(x)dx = \int f[g(t)]g'(t)dt, \text{ where } x = g(t).$$

(b) Integration by Partial Fractions:

Consider $\frac{f(x)}{g(x)}$ defines a rational polynomial function.

➤ In rational polynomial function if the degree (i.e., highest power of the variable) of numerator (Nr.) is **greater than or equal to** the degree of denominator (Dr.), then (without any doubt) **always perform the division** i.e., divide the Nr. by Dr. before doing anything and thereafter use the following:

$$\frac{\text{Numerator}}{\text{Denominator}} = \text{Quotient} + \frac{\text{Remainder}}{\text{Denominator}}$$

Table Demonstrating Partial Fractions or Various Forms

Form of the Rational Function	Form of the Partial Fraction
$\frac{px+q}{(x-a)(x-b)}, a \neq b$	$\frac{A}{x-a} + \frac{B}{x-b}$
$\frac{px+q}{(x-a)^2}$	$\frac{A}{x-a} + \frac{B}{(x-a)^2}$
$\frac{px^2+qx+r}{(x-a)(x-b)(x-c)}$	$\frac{A}{x-a} + \frac{B}{x-b} + \frac{C}{x-c}$
$\frac{px^2+qx+r}{(x-a)^2(x-b)}$	$\frac{A}{x-a} + \frac{B}{(x-a)^2} + \frac{C}{x-b}$

$$\frac{px^2 + qx + r}{(x-a)(x^2 + bx + c)}$$

where $x^2 + bx + c$ can't be factorized further.

$$\frac{A}{x-a} + \frac{Bx+C}{x^2+bx+c}$$

(c) Integration by Parts :

If U and V be two functions of x , then

$$\int U_{(i)} V_{(ii)} dx = U \int V dx - \int \left\{ \frac{dU}{dx} \int V dx \right\} dx$$

Key Formulae

Formulae for Indefinite Integrals

(a) $\int x^n dx = \frac{x^{n+1}}{n+1} + C, n \neq -1$

(b) $\int \frac{1}{x} dx = \log |x| + C$

(c) $\int a^x dx = \frac{1}{\log a} a^x + C$

(d) $\int e^{ax} dx = \frac{1}{a} e^{ax} + C$

(e) $\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + C$

(f) $\int \cos(ax) dx = \frac{1}{a} \sin(ax) + C$

(g) $\int \tan x dx = \log |\sec x| + C$ or $-\log |\cos x| + C$

(h) $\int \cot x dx = \log |\sin x| + C$ or $-\log |\operatorname{cosec} x| + C$

(i) $\int \sec x dx = \log |\sec x + \tan x| + C$ or $\log \left| \tan \left(\frac{\pi}{4} + \frac{x}{2} \right) \right| + C$

(j) $\int \operatorname{cosec} x dx = \log |\operatorname{cosec} x - \cot x| + C$ or $\log \left| \tan \frac{x}{2} \right| + C$

(k) $\int \sec^2 x dx = \tan x + C$

(l) $\int \operatorname{cosec}^2 x dx = -\cot x + C$

(m) $\int \sec x \cdot \tan x dx = \sec x + C$

(n) $\int \operatorname{cosec} x \cdot \cot x dx = -\operatorname{cosec} x + C$

(o) $\int \frac{1}{x\sqrt{x^2-1}} dx = \sec^{-1} x + C$

(p) $\int \frac{1}{a^2+x^2} dx = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + C$

(q) $\int \frac{1}{a^2-x^2} dx = \frac{1}{2a} \log \left| \frac{a+x}{a-x} \right| + C$

(r) $\int \frac{1}{x^2-a^2} dx = \frac{1}{2a} \log \left| \frac{x-a}{x+a} \right| + C$

(s) $\int \frac{1}{\sqrt{x^2-a^2}} dx = \log \left| x + \sqrt{x^2-a^2} \right| + C$

(t) $\int \frac{1}{\sqrt{x^2+a^2}} dx = \log \left| x + \sqrt{x^2+a^2} \right| + C$

(u) $\int \frac{1}{\sqrt{a^2-x^2}} dx = \sin^{-1} \left(\frac{x}{a} \right) + C$

(v) $\int \frac{1}{ax+b} dx = \frac{1}{a} \log |ax+b| + C$

(w) $\int \lambda dx = \lambda x + C$, where ' λ ' is a constant.

(x) $\int \sqrt{x^2-a^2} dx = \frac{x}{2} \sqrt{x^2-a^2} - \frac{a^2}{2} \log \left| x + \sqrt{x^2-a^2} \right| + C$

(y) $\int \sqrt{x^2+a^2} dx = \frac{x}{2} \sqrt{x^2+a^2} + \frac{a^2}{2} \log \left| x + \sqrt{x^2+a^2} \right| + C$

(z) $\int \sqrt{a^2-x^2} dx = \frac{x}{2} \sqrt{a^2-x^2} + \frac{a^2}{2} \sin^{-1} \left(\frac{x}{a} \right) + C$

Topic-2 Definite Integral

Concepts Covered • Second fundamental theorem, • Properties of definite integral.



Revision Notes

➤ **Meaning of Definite Integral of Function**

If $\int f(x)dx = F(x)$ i.e., $F(x)$, be an integral of $f(x)$, then $F(b) - F(a)$ is called the definite integral of $f(x)$ between the limits a and b and in symbols it is written as $\int_a^b f(x)dx =$

$[F(x)]_a^b$. Moreover, the definite integral gives a unique and definite value (numeric value) of **anti-derivative** of the function between the given intervals. It acts as a substitute for evaluating the area analytically.

Key Formulae

(a) $\int_a^b f(x)dx = F(b) - F(a)$

(b) $\int_a^b f(x)dx = -\int_b^a f(x)dx$

(c) $\int_a^b f(x)dx = \int_a^b f(t)dt \quad (dx = dt)$

(d) $\int_a^b f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx, a < c < b$

(e) $\int_0^a f(x)dx = \int_0^a f(a-x)dx$

(f) $\int_a^b f(x)dx = \int_a^b f(a+b-x)dx$

(g) $\int_{-a}^a f(x)dx = \begin{cases} 2\int_0^a f(x)dx, & \text{if } f(x) \text{ is an even function i.e., } f(-x) = f(x) \\ 0, & \text{if } f(x) \text{ is an odd function i.e., } f(-x) = -f(x) \end{cases}$

(h) $\int_{-a}^a f(x)dx = \int_0^a \{f(x) + f(-x)\}dx$


(i) $\int_0^{2a} f(x)dx = \begin{cases} 2\int_0^a f(x)dx, & \text{if } f(2a-x) = f(x) \\ 0 & ; \text{ if } f(2a-x) = -f(x) \end{cases}$

(j) $\int_0^{2a} f(x)dx = \int_0^{2a} \{f(x) + f(2a-x)\}dx$




Mnemonics

SeCond FundAmental Theorem of Definite Integration



You can also remember

fcc is small **f**ashionable **C**lothes



Interpretation :

Let f be a continuous function defined on a closed interval $[a, b]$ and F be an anti derivative of f . Then

$\int_a^b f(x)dx = [F(x)]_a^b = F(b) - F(a)$, where a and b are called limit of Integration.

CHAPTER-8

APPLICATIONS OF THE INTEGRALS



Revision Notes

➤ Area Under Simple Curves:


- (i) Let us find the area bounded by the curve $y = f(x)$, X-axis and the ordinates $x = a$ and $x = b$. Consider the area under the curve as composed by large number of thin vertical strips.

Let there be an **arbitrary** strip of height y and width dx .

Area of elementary strip $dA = y dx$, where $y = f(x)$.

Total area A of the region between X-axis ordinates $x = a$, $x = b$ and the curve $y = f(x) =$ sum of areas of elementary thin strips across the region PQML.

$$A = \int_a^b y dx = \int_a^b f(x) dx$$



Key Word


Arbitrary: In mathematics, "arbitrary" just means "for all".

For example: "For all a, b , $a + b = b + a$ ".

Another way to say this would be " $a + b = b + a$ for arbitrary a, b ."


- (ii) The area A of the region bounded by the curve $x = g(y)$, Y-axis and the lines $y = c$ and $y = d$ is given by

$$A = \int_c^d x dy = \int_c^d g(y) dy$$



- (iii) If the curve under consideration lies below X-axis, then $f(x) < 0$ from $x = a$ to $x = b$, the area bounded by the curve $y = f(x)$ and the ordinates $x = a$, $x = b$ and X-axis is negative. But, if the numerical value of the area is to be taken into consideration, then

$$\text{Area} = \left| \int_a^b f(x) dx \right|$$



- (iv) It may also happen that some portion of the curve is above X-axis and some portion is below X-axis as shown in the figure. Let A_1 be the area below X-axis and A_2 be the area above the X-axis. Therefore, area bounded by the curve $y = f(x)$, X-axis and the ordinates $x = a$ and $x = b$ is given by

$$A = |A_1| + |A_2|$$

CHAPTER-9

DIFFERENTIAL EQUATIONS

Topic-1

Basic Differential Equations

Concepts Covered • Order of differential Equation

• Degree of differential Equation



Revision Notes

Orders and Degrees of Differential Equation:

- We shall prefer to use the following notations for derivatives.
- $\frac{dy}{dx} = y'$, $\frac{d^2y}{dx^2} = y''$, $\frac{d^3y}{dx^3} = y'''$
- For derivatives of higher order, it will be in convenient to use so many dashes as super suffix therefore, we use the notation y_n for n^{th} order derivative $\frac{d^n y}{dx^n}$.
- Order and degree (if defined) of a **differential equation** are always positive integers.



Key Word

Differential Equation: In Mathematics, a differential equation is an equation with one or more derivatives of a function. The derivative of the function is given by dy/dx . In other words, it is defined as the equation that contains derivatives of one or more dependent variables with respect to one or more independent variables.



Know the terms

- **Order of a differential equation:** It is the order of the highest order derivative appearing in the differential equation.
- **Degree of a differential equation:** It is the degree (power) of the highest order derivative, when the differential coefficients are made free from the radicals and the fractions.

Topic-2

Variable Separable Methods

Concepts Covered • General Solution, • Particular Solutions, • Variable Separable Method



Revision Notes

Solutions of a differential equation:

(a) **General Solution:** The solution which contains as many as arbitrary constants as the order of the differential equations, e.g., $y = \alpha \cos x + \beta \sin x$ is the general solution of $\frac{d^2y}{dx^2} + y = 0$.

(b) **Particular Solution:** Solution obtained by giving particular values to the arbitrary constants in the general solution of a differential equation is

called a particular solution e.g., $y = 3 \cos x + 2 \sin x$ is a particular solution of the differential equation

$$\frac{d^2y}{dx^2} + y = 0.$$

(c) **Solution of Differential by Variable Separable Method:** A **variable** separable form of the differential equation is the one which can be expressed in the form of $f(x)dx = g(y)dy$. The solution is given by $\int f(x)dx = \int g(y)dy + k$ where k is the **constant** of integration.

Topic-3**Linear Differential Equations****Concepts Covered** • Linear Differential Equations in x only and in y only**Revision Notes**

➤ **Linear differential equation in y :** It is of the form $\frac{dy}{dx} + P(x)y = Q(x)$, where $P(x)$ and $Q(x)$ are

functions of x only.

➤ **Solving Linear Differential Equation in y :**

STEP 1: Write the given differential equation in the form

$$\frac{dy}{dx} + P(x)y = Q(x).$$

STEP 2: Find the Integration Factor (I.F.) = $e^{\int P(x)dx}$.

STEP 3: The solution is given by, $y.(I.F.) = \int Q(x).I.F. + k$

where k is the constant of integration.

➤ **Linear differential equation in x :** It is of the form $\frac{dx}{dy} + P(y)x = Q(y)$, where $P(y)$ and $Q(y)$

are functions of y only.


➤ **Solving Linear Differential Equation in x :**

STEP 1: Write the given differential equation in the form

$$\frac{dx}{dy} + P(y)x = Q(y).$$

STEP 2: Find the Integration Factor (I.F.) = $e^{\int P(y)dy}$.

STEP 3: The solution is given by, $x.(I.F.) = \int Q(y).(I.F.)dy + \lambda$, where λ is the constant of integration.

**Mnemonics****Linear Differential Equations****SOLDE-YIF-EIQ-IFC**

Son Of Liladhar Dixit Eklavya (SOLDE)

& WIFe (Y I F) Exploring Indian Qilla

for **International Fitness Certificate**

$$y. IF = \int Q. IF + C$$

S — Solution

D — Differential

O — Of

E — Equation

L — Linear

Interpretation:

Differential equation is of the form $\frac{dy}{dx} + py = Q$, where

P and Q are constants or the function of ' x ' is called a first order linear differential equations. Its solution is

given as $Y.IF = \int Q.IF + C$

Topic-4

Homogeneous Differential Equations

Concepts Covered • Solution of Homogenous Differential Equation of first order and first degree



Revision Notes

➤ **Homogeneous Differential Equations and their solution:**

➤ **Identifying a Homogeneous Differential equation:**

STEP 1: Write down the given differential equation in the form $\frac{dy}{dx} = f(x, y)$.

STEP 2: If $f(kx, ky) = k^n f(x, y)$, then the given differential equation is **homogeneous** of degree 'n'.

➤ **Solving a Homogeneous Differential Equation:**

CASE I: If $\frac{dy}{dx} = f(x, y)$

Put $y = vx$
 or $\frac{dy}{dx} = v + x \frac{dv}{dx}$

CASE II: If $\frac{dx}{dy} = f(x, y)$


Put $x = vy$
 or $\frac{dx}{dy} = v + y \frac{dv}{dy}$

Then, we separate the variables to get the required solution.



Mnemonics

Homogeneous Differential Equation



Interpretation:


Differential equation can be expressed in the form $\frac{dy}{dx} = f(x, y)$ or $\frac{dx}{dy} = g(x, y)$ where $f(x, y)$ and $g(x, y)$ are homogeneous

functions of sum is called a homogeneous Differential equation. These equations can be solved by substituting $y = vx$ so that dependent variable y is changed to another variable v , where v is some unknown function.

UNIT – IV: VECTORS & THREE-DIMENSIONAL GEOMETRY**CHAPTER-10****VECTORS****Basic Algebra of Vectors****Topic-1**

Concepts Covered • Basic concepts of vectors,
• Operations on vectors • Different types of vectors,
• Triangle Law, • Parallelogram Law

**Revision Notes****1. Vector: Basic Introduction:**

- A physical quantity having **magnitude** as well as the direction is called a vector. It is denoted as \vec{AB} or \vec{a} . Its magnitude (or modulus) is $|\vec{AB}|$ or $|\vec{a}|$ otherwise, simply AB or a .
- Vectors are denoted by symbols such as \vec{a} .

 [Pictorial representation of vector]

2. Initial and Terminal Points:

The initial and terminal points means that point from which the vector originates and terminates respectively.

**Key Words**

Magnitude: It is defined as the maximum extent of size and the direction of an object. Magnitude is used as a common factor in vector and scalar quantities.

3. Position Vector:

The position vector of a point say $P(x, y, z)$ is $\vec{OP} = \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$ and the magnitude is $|\vec{r}| = \sqrt{x^2 + y^2 + z^2}$.

- Also, $\vec{AB} = (\text{Position Vector of } B) - (\text{Position Vector of } A)$. For example, let $A(x_1, y_1, z_1)$ and $B(x_2, y_2, z_2)$. Then, $\vec{AB} = (x_2\hat{i} + y_2\hat{j} + z_2\hat{k}) - (x_1\hat{i} + y_1\hat{j} + z_1\hat{k})$.
- Here, \hat{i} , \hat{j} and \hat{k} are the unit vectors along the axes OX , OY and OZ respectively (The


discussion about unit vectors is given later under 'types of vectors').

4. Addition of vectors


- (a) **Triangular law:** If two adjacent sides (say sides AB and BC) of a triangle ABC are represented by \vec{a} and \vec{b} taken in same order, then the third side of the triangle taken in the reverse order gives the sum of vectors \vec{a} and \vec{b} i.e.,

$$\vec{AC} = \vec{AB} + \vec{BC} \Rightarrow \vec{AC} = \vec{a} + \vec{b}.$$

- Also since $\vec{AC} = -\vec{CA} \Rightarrow \vec{AB} + \vec{BC} + \vec{CA} = \vec{0}$.



- (b) **Parallelogram law:** If two vectors \vec{a} and \vec{b} are represented in magnitude and the direction by the two adjacent sides (say OA and OB) of a parallelogram $OACB$, then their sum is given by that diagonal of parallelogram which is co-initial with \vec{a} and \vec{b} i.e., $\vec{OC} = \vec{OA} + \vec{OB}$.



Note: Multiplication of a vector by a scalar

Let \vec{a} be any vector and k be any non-zero scalar. Then the product $k\vec{a}$ is defined as a vector whose magnitude is $|k|$ times that of \vec{a} and the direction is

- (i) same as that of \vec{a} if k is positive, and
 (ii) opposite as that of \vec{a} if k is negative.

**Key Fact**

- Vector calculus and its sub objective vector fields was invented by two men J. Willard Gibbs and Oliver Heaviside at the end of 19th century.

**Know the Terms****Types of Vectors:**

- (a) **Zero or Null vector:** It is that vector whose initial and terminal points are coincident. It is denoted by $\vec{0}$. Of course its magnitude is 0 (zero).
- Any non-zero vector is called a **proper vector**.
- (b) **Co-initial vectors:** Those vectors (two or more) having the same starting point are called the co-initial vectors.
- (c) **Co-terminus vectors:** Those vectors (two or more) having the same terminal point are called the co-terminus vectors.
- (d) **Negative of a vector:** The vector which has the same magnitude as the \vec{r} but opposite direction. It is denoted by $-\vec{r}$. Hence if,
- $\vec{AB} = \vec{r}$ or $\vec{BA} = -\vec{r}$ i.e., $\vec{AB} = -\vec{BA}$, $\vec{PQ} = -\vec{QP}$ etc.
- (e) **Unit vector:** It is a vector with the unit magnitude.

The unit vector in the direction of vector \vec{r} is given by $\hat{r} = \frac{\vec{r}}{|\vec{r}|}$ such that $|\hat{r}| = 1$, so, if

$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$ then its unit vector is:

$$\hat{r} = \frac{x}{\sqrt{x^2 + y^2 + z^2}}\hat{i} + \frac{y}{\sqrt{x^2 + y^2 + z^2}}\hat{j} + \frac{z}{\sqrt{x^2 + y^2 + z^2}}\hat{k}.$$

- Unit vector perpendicular to the plane \vec{a} and

$$\vec{b} \text{ is: } \pm \frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|}.$$

- (f) **Reciprocal of a vector:** It is a vector which has the same direction as the vector \vec{r} but magnitude

equal to the reciprocal of the magnitude of \vec{r} .

It is denoted as \vec{r}^{-1} . Hence $|\vec{r}^{-1}| = \frac{1}{|\vec{r}|}$.

- (g) **Equal vectors:** Two vectors are said to be equal if they have the same magnitude as well as direction, regardless of the position of their initial points.

Thus $\vec{a} = \vec{b} \Leftrightarrow \begin{cases} |\vec{a}| = |\vec{b}| \\ \vec{a} \text{ and } \vec{b} \text{ have same direction} \end{cases}$

Also, if $\vec{a} = \vec{b} \Rightarrow a_1\hat{i} + a_2\hat{j} + a_3\hat{k} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$

$$\Rightarrow a_1 = b_1, a_2 = b_2, a_3 = b_3.$$

- (h) **Collinear or Parallel vector:** Two vectors \vec{a} and \vec{b} are collinear or parallel if there exists a non-zero scalar λ such that $\vec{a} = \lambda\vec{b}$.
- It is important to note that the respective coefficients of $\hat{i}, \hat{j}, \hat{k}$ in \vec{a} and \vec{b} are proportional provided they are parallel or collinear to each other.
 - The d.r.'s of parallel vectors are same (or are in proportion).
 - The vectors \vec{a} and \vec{b} will have same or opposite direction as λ is positive or negative respectively.
 - The vectors \vec{a} and \vec{b} are collinear if $\vec{a} \times \vec{b} = \vec{0}$.
- (i) **Free vectors:** The vectors which can undergo parallel displacement without changing its magnitude and direction are called free vectors.

Key Formulæ

The position vector of a point say P dividing a line segment joining the points A and B whose position vectors are \vec{a} and \vec{b} respectively, in the ratio $m : n$.

(a) Internally, $\vec{OP} = \frac{m\vec{b} + n\vec{a}}{m+n}$


(b) Externally, $\vec{OP} = \frac{m\vec{b} - n\vec{a}}{m-n}$

- Also if point P is the mid-point of line segment AB , then $\vec{OP} = \frac{\vec{a} + \vec{b}}{2}$.



Mnemonics

Types Of Vectors (A)




Interpretation :

Types of Vectors-

- Zero Vector** - Initial and terminal points coincide
- Unit Vector** - Magnitude is unity
- Coinitial Vectors** - Same initial points
- Collinear vectors** - Parallel to the same Line
- Equal Vectors** - Same magnitude and direction
- Negative of a vector** - Same magnitude, opp. direction

Properties Of Vectors(B)


"Neither choose East nor choose north, always choose North-East and save your time".



Mnemonics

Interpretation :


The vector sum of two co-initial vectors is given by the diagonal of the parallelogram whose adjacent sides are given vectors.



$$\vec{AB} + \vec{AC} = \vec{AD}$$

Properties Of Vectors(C)

Aao Bnaye Circle Tirahe par



Interpretation:

The vector sum of the three sides of a triangle taken in order is $\vec{0}$ i.e

$$\vec{AB} + \vec{BC} + \vec{CA} = \vec{0}$$

Topic-2

Dot Product of Vectors

Concepts Covered • Properties of dot product, • Projection of a vector




Revision Notes

Scalar Product or Dot Product: The dot product of two vectors \vec{a} and \vec{b} is defined by,

$$\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta \text{ where } \theta \text{ is the angle between}$$

$$\vec{a} \text{ and } \vec{b}, 0 \leq \theta \leq \pi.$$



Consider

$$\vec{a} = a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}, \vec{b} = b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k},$$

then $\vec{a} \cdot \vec{b} = a_1 b_1 + a_2 b_2 + a_3 b_3$.

⇒ **Projection of a vector :** \vec{a} on the other vector say \vec{b} is given as $\left(\frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} \right)$.

⇒ **Projection of a vector :** \vec{b} on the other vector say \vec{a} is given as $\left(\frac{\vec{a} \cdot \vec{b}}{|\vec{a}|} \right)$.

Key Word

Projection: The image of a geometrical figure reproduced on a line, plane or surface.



Know the Properties (Dot Product)

• **Properties/Observations of Dot product**

⇒ $\hat{i} \cdot \hat{i} = |\hat{i}| |\hat{i}| \cos 0 = 1$ or $\hat{i} \cdot \hat{i} = 1 = \hat{j} \cdot \hat{j} = \hat{k} \cdot \hat{k}$

⇒ $\hat{i} \cdot \hat{j} = |\hat{i}| |\hat{j}| \cos \frac{\pi}{2} = 0$ or $\hat{i} \cdot \hat{j} = 0 = \hat{j} \cdot \hat{k} = \hat{k} \cdot \hat{i}$

⇒ $\vec{a} \cdot \vec{b} \in R$, where R is real number i.e., any scalar.

⇒ $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{a}$ (Commutative property of dot product).

⇒ $\vec{a} \cdot \vec{b} = 0 \Leftrightarrow \vec{a} \perp \vec{b}$ or $|\vec{a}| = 0$ or $|\vec{b}| = 0$.

⇒ If $\theta = 0$, then $\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}|$. Also

$\vec{a} \cdot \vec{a} = |\vec{a}|^2 = a^2$; as θ in this case is 0.

Moreover if $\theta = \pi$, then $\vec{a} \cdot \vec{b} = -|\vec{a}| |\vec{b}|$.

⇒ $\vec{a} \cdot (\vec{b} + \vec{c}) = \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c}$

(Distributive property of dot product).

⇒ $\vec{a} \cdot (-\vec{b}) = -(\vec{a} \cdot \vec{b}) = (-\vec{a}) \cdot \vec{b}$.



Key Formulae

⇒ Angle between two vectors \vec{a} and \vec{b} can be found by the expression given below :

$$\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} \text{ or } \theta = \cos^{-1} \left(\frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} \right)$$

Topic-3

Cross Product

Concepts Covered • Properties of cross product, • Relationship between Vector product and scalar product




Revision Notes

1. The **cross product** or vector product of two vectors \vec{a} and \vec{b} is defined by,

$$\vec{a} \times \vec{b} = |\vec{a}| |\vec{b}| \sin \theta \hat{n}, \text{ where } \theta \text{ is the angle}$$

between the vectors \vec{a} and \vec{b} , $0 \leq \theta \leq \pi$ and \hat{n} is a unit vector perpendicular to both \vec{a} and \vec{b} .

For better illustration, see figure.



Consider $\vec{a} = a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}$, $\vec{b} = b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k}$.

then

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} =$$

$$(a_2 b_3 - a_3 b_2) \hat{i} - (a_1 b_3 - a_3 b_1) \hat{j} + (a_1 b_2 - a_2 b_1) \hat{k}.$$

• **Properties/Observations of Cross Product**

$$\Rightarrow \hat{i} \times \hat{i} = \hat{j} \times \hat{j} = \hat{k} \times \hat{k} = \vec{0} \text{ or } \hat{i} \times \hat{i} = \vec{0} = \hat{j} \times \hat{j} = \hat{k} \times \hat{k}.$$

$$\Rightarrow \hat{i} \times \hat{j} = \hat{j} \times \hat{k} = \hat{k} \times \hat{i} = \hat{i} \text{ and } \hat{j} \times \hat{i} = \hat{i} \times \hat{j} = -\hat{k} \text{ or } \hat{i} \times \hat{j} = \hat{k}, \hat{j} \times \hat{k} = \hat{i}, \hat{k} \times \hat{i} = \hat{j}.$$

$\Rightarrow \vec{a} \times \vec{b}$ is a vector \vec{c} (say) then this vector \vec{c} is perpendicular to both the vectors \vec{a} and \vec{b} .

$$\Rightarrow \vec{a} \times \vec{b} = \vec{0} \Leftrightarrow \vec{a} \parallel \vec{b} \text{ or } \vec{a} = \vec{0}, \vec{b} = \vec{0}.$$

$$\Rightarrow \vec{a} \times \vec{a} = \vec{0}.$$

$\Rightarrow \vec{a} \times \vec{b} \neq \vec{b} \times \vec{a}$ (Commutative property does not hold for cross product).

$$\Rightarrow \vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c} \text{ (Left distributive).}$$

$$\Rightarrow (\vec{b} + \vec{c}) \times \vec{a} = \vec{b} \times \vec{a} + \vec{c} \times \vec{a} \text{ (Right distributive).}$$

(Distributive property of the vector product or cross product)

2. Relationship between Vector product and Scalar product [Lagrange's Identity]

$$\text{or } |\vec{a} \times \vec{b}|^2 + (\vec{a} \cdot \vec{b})^2 = |\vec{a}|^2 \cdot |\vec{b}|^2$$

3. Cauchy-Schwarz inequality :

For any two vectors \vec{a} and \vec{b} , always have $|\vec{a} \cdot \vec{b}| \leq |\vec{a}| |\vec{b}|$.

Note:

- If \vec{a} and \vec{b} represent the adjacent sides of a triangle, then the area of triangle can be obtained by evaluating $\frac{1}{2} |\vec{a} \times \vec{b}|$.
- If \vec{a} and \vec{b} represent the adjacent sides of a parallelogram, then the area of parallelogram can be obtained by evaluating $|\vec{a} \times \vec{b}|$.
- The area of the parallelogram with diagonals \vec{a} and \vec{b} is $\frac{1}{2} |\vec{a} \times \vec{b}|$.



Key Formulae

\Rightarrow Angle between two vectors \vec{a} and \vec{b} in terms of cross-product can be found by the expression given here :

$$\sin \theta = \frac{|\vec{a} \times \vec{b}|}{|\vec{a}| |\vec{b}|} \text{ or } \theta = \sin^{-1} \left(\frac{|\vec{a} \times \vec{b}|}{|\vec{a}| |\vec{b}|} \right)$$

CHAPTER-11

THREE DIMENSIONAL GEOMETRY

Topic-1

Direction Ratios and Direction Cosines

Concepts Covered • Direction Ratios, • Direction Cosines

• Relationship between DC's of a line.



Revision Notes

1. Direction Cosines of a Line:

- If A and B are two points on a given line L, then direction cosines of vectors

\vec{AB} and \vec{BA} are the direction cosines (d.c.'s) of line L. Thus if α, β, γ are the direction-angles which the line L makes with the positive

direction of X, Y, Z-axis respectively, then its d.c.'s are $\cos \alpha, \cos \beta, \cos \gamma$.

- If direction of line L is reversed, the direction angles are replaced by their **supplements angles** i.e., $\pi - \alpha, \pi - \beta, \pi - \gamma$ and so are the d.c.'s i.e., the direction cosines become $-\cos \alpha, -\cos \beta, -\cos \gamma$.



Key Word

Supplement angles: Two angles or arcs whose sum is 180° degrees.

- So, a line in space has two set of d.c.'s viz $\pm \cos \alpha, \pm \cos \beta, \pm \cos \gamma$.
- The d.c.'s are generally denoted by l, m, n . Also $l^2 + m^2 + n^2 = 1$ (relation between Direction Cosines) and so we can deduce that $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$. Also, $\sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma = 2$.
- The d.c.'s of a line joining the points $A(x_1, y_1, z_1)$ and $B(x_2, y_2, z_2)$ are $\pm \frac{x_2 - x_1}{AB}, \pm \frac{y_2 - y_1}{AB}, \pm \frac{z_2 - z_1}{AB}$;

where AB is the distance between the points A and B i.e.,

$$AB = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

2. Direction Ratios of a Line:

Any three numbers a, b, c (say) which are proportional to d.c.'s i.e., l, m, n of a line are called the **direction ratios** (d.r.'s) of the line. Thus, $a = \lambda l, b = \lambda m, c = \lambda n$ for any $\lambda \in \mathbb{R} - \{0\}$.

Consider, $\frac{l}{a} = \frac{m}{b} = \frac{n}{c} = \frac{1}{\lambda}$ (say)

or $l = \frac{a}{\lambda}, m = \frac{b}{\lambda}, n = \frac{c}{\lambda}$

or $\left(\frac{a}{\lambda}\right)^2 + \left(\frac{b}{\lambda}\right)^2 + \left(\frac{c}{\lambda}\right)^2 = 1$ [Using $l^2 + m^2 + n^2 = 1$]

or $\lambda = \pm \sqrt{a^2 + b^2 + c^2}$

Therefore, $l = \pm \frac{a}{\sqrt{a^2 + b^2 + c^2}},$
 $m = \pm \frac{b}{\sqrt{a^2 + b^2 + c^2}}, n = \pm \frac{c}{\sqrt{a^2 + b^2 + c^2}}$

- The d.r.'s of a line joining the points $A(x_1, y_1, z_1)$ and $B(x_2, y_2, z_2)$ are $x_2 - x_1, y_2 - y_1, z_2 - z_1$ or $x_1 - x_2, y_1 - y_2, z_1 - z_2$.
- Direction ratios are sometimes called as **Direction Numbers**.



Key Formulæ

1. Distance Formula:

The distance between two points $A(x_1, y_1, z_1)$ and $B(x_2, y_2, z_2)$ is given by the expression

$$AB = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2} \text{ units.}$$

2. Section Formula:

The co-ordinates of a point Q which divides the line joining the points $A(x_1, y_1, z_1)$ and $B(x_2, y_2, z_2)$ in the ratio $m : n$

(a) internally, are $\left(\frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n}, \frac{mz_2 + nz_1}{m+n} \right)$

(b) externally, are $\left(\frac{mx_2 - nx_1}{m-n}, \frac{my_2 - ny_1}{m-n}, \frac{mz_2 - nz_1}{m-n} \right)$.



Amazing Facts

- The largest 3D shape in the world is a Rhombicosidodecahedron. It is an Archimedean solid. It has 20 faces that are triangular, 30 faces that are squares, and 12 are that are pentagons. This shape has 120 edges and 60 vertices.
- The Louvre pyramid is a beautiful installation that is perfect example of a 3D shape i.e., square pyramid. It is situated in the city of Paris in the prestigious museum of the Louvre.



Mnemonics


Direction Cosines

1 glass L e M o N juice

$$l^2 + m^2 + n^2 = 1$$

Interpretation:

Direction cosines of a line are the cosines of the angles made by the line with the positive directions of the co. ordinate axes. If l, m, n are the d. c. s of a line, then $l^2 + m^2 + n^2 = 1$

Dance Choreographer Prefer Dieting**Direction Ratios****Director Remo a Professional Dancer****Choreographer created****3 Lifetime Movies with New faces a b c**

$$l = \frac{a}{\sqrt{a^2+b^2+c^2}}, m = \frac{b}{\sqrt{a^2+b^2+c^2}}, n = \frac{c}{\sqrt{a^2+b^2+c^2}}$$

Lines & Its Equations in Different forms**Topic-2****Concepts Covered** • Equation of line in cartesian and vector form,

- Shortest distance between lines
- Skew lines
- Condition of parallelism and perpendicularity of lines.

**Revision Notes****1. Equation of a Line passing through two given points:**

Consider the two given points as $A(x_1, y_1, z_1)$ and $B(x_2, y_2, z_2)$ with position vectors \vec{a} and \vec{b} respectively. Also assume \vec{r} as the position vector of any arbitrary point $P(x, y, z)$ on the line L passing through A and B . Thus,

$$\vec{OA} = \vec{a} = x_1 \hat{i} + y_1 \hat{j} + z_1 \hat{k}, \vec{OB} = \vec{b} = x_2 \hat{i} + y_2 \hat{j} + z_2 \hat{k},$$

$$\vec{OP} = \vec{r} = x \hat{i} + y \hat{j} + z \hat{k}$$

- (a) **Vector equation of a line:** Since, the points A, B and P all lie on the same line which means that they are all collinear points.

Further it means, $\vec{AP} = \vec{r} - \vec{a}$ and $\vec{AB} = \vec{b} - \vec{a}$ are collinear vectors, i.e.,

$$\vec{AP} = \lambda \vec{AB}$$

$$\text{or } \vec{r} - \vec{a} = \lambda(\vec{b} - \vec{a})$$

$$\text{or } \vec{r} = \vec{a} + \lambda(\vec{b} - \vec{a}), \text{ where } \lambda \in R.$$

This is the vector equation of the line.

- (b) **Cartesian equation of a line:** By using the vector equation of the line $\vec{r} = \vec{a} + \lambda(\vec{b} - \vec{a})$, we get

$$x \hat{i} + y \hat{j} + z \hat{k} = x_1 \hat{i} + y_1 \hat{j} + z_1 \hat{k} +$$

$$\lambda [(x_2 - x_1) \hat{i} + (y_2 - y_1) \hat{j} + (z_2 - z_1) \hat{k}]$$

On equating the coefficients of $\hat{i}, \hat{j}, \hat{k}$, we get

$$x = x_1 + \lambda(x_2 - x_1), y = y_1 + \lambda(y_2 - y_1),$$

$$z = z_1 + \lambda(z_2 - z_1) \quad \dots(i)$$

On eliminating λ , we have

$$\frac{x - x_1}{x_2 - x_1} = \frac{y - y_1}{y_2 - y_1} = \frac{z - z_1}{z_2 - z_1}$$

2. Angle between two lines:

- (a) **When d.r.'s or d.c.'s of the two lines are given:**

Consider two lines L_1 and L_2 with d.r.'s in proportion to a_1, b_1, c_1 and a_2, b_2, c_2 respectively; d.c.'s as l_1, m_1, n_1 and l_2, m_2, n_2 .

$$\text{Then, } \cos \theta = \frac{a_1 a_2 + b_1 b_2 + c_1 c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}}$$

- Also, in terms of d.c.'s: $\cos \theta = |l_1 l_2 + m_1 m_2 + n_1 n_2|$.
- Sine of angle is given as:

$$\sin \theta = \frac{\sqrt{(a_1 b_2 - a_2 b_1)^2 + (b_1 c_2 - b_2 c_1)^2 + (c_1 a_2 - c_2 a_1)^2}}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}}$$

(b) When Vector equations of two lines are given:

Consider vector equations of lines L_1 and L_2 as

$$\vec{r}_1 = \vec{a}_1 + \lambda \vec{b}_1 \text{ and } \vec{r}_2 = \vec{a}_2 + \mu \vec{b}_2 \text{ respectively.}$$

Then, the acute angle θ between the two lines is given by the relation

$$\cos \theta = \frac{|\vec{b}_1 \cdot \vec{b}_2|}{\|\vec{b}_1\| \|\vec{b}_2\|}$$

(c) When Cartesian equation of two lines are given:

Consider the lines L_1 and L_2 in Cartesian form as,

$$L_1: \frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1}$$

$$L_2: \frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}$$

Then the acute angle θ between the lines L_1 and L_2 can be obtained by,

$$\cos \theta = \frac{|a_1 a_2 + b_1 b_2 + c_1 c_2|}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}}$$

Note:

• For two perpendicular lines : $a_1 a_2 + b_1 b_2 + c_1 c_2 = 0$, $l_1 l_2 + m_1 m_2 + n_1 n_2 = 0$.

• For two parallel lines :

$$\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}; \quad \frac{l_1}{l_2} = \frac{m_1}{m_2} = \frac{n_1}{n_2}$$

3. Shortest Distance between Two Lines:

If two lines are in the same plane i.e., they are coplanar, they will intersect each other if they are non-parallel. Hence, the shortest distance between them is zero. "If the lines are parallel then the shortest distance between them will be the perpendicular distance between the lines i.e., the length of the perpendicular drawn from a point on one line onto the other line". Adding to this discussion, in space, there are lines which are neither intersecting nor parallel. In fact, such pair of lines are non-coplanar and are called the skew lines.

UNIT – V: LINEAR PROGRAMMING**CHAPTER-12****LINEAR PROGRAMMING****Revision Notes**

Linear programming problems: Problems which minimize or maximize a linear function z subject to certain conditions determined by a set of linear inequalities with non-negative variables are known as linear programming problems.

Objective function: A linear function $z = ax + by$, where a and b are constants which has to be maximised or minimised according to a set of given

conditions, is called as linear objective function.

Decision variables: In the objective function $z = ax + by$, the variables x, y are said to be decision variables.

Constraints: The restrictions in the form of inequalities on the variables of a linear programming problems are called constraints. The condition $x \geq 0, y \geq 0$ are known as non-negative restrictions.

**Key Terms**

Feasible region: The common region determined by all the constraints including non-negative constraints $x, y \geq 0$ of linear programming problem is known as the feasible region.

Feasible solution: Points with in and on the boundary of the feasible region represents feasible solutions of constraints.

In the feasible region, there are infinitely many points

(solutions) which satisfy the given conditions.

Theorem 1: Let R be the feasible region for a linear programming problem and let $Z = ax + by$ be the objective function. When Z has an optimal value (maximum or minimum), where variables x and y are subject to constraints described by linear inequalities, the optimal value must occur at a corner point (vertex) of the feasible region.

Theorem 2: Let R be the feasible region for a linear programming problem, and let $Z = ax + by$ be the objective function. If R is bounded, then the objective function Z has both maximum and minimum values of R and each of these occurs at a corner point (vertex) of R .

However, if the feasible region is unbounded, the optimal value obtained may not be maximum or minimum.



Mnemonics

LLP parameters

N	O	C
↓	↓	↓
Non-negative variables	Objective function	Constraints



Key Facts

- Linear programming is often used for problems where no exact solution is known, for example for planning traffic flows.
- Linear programming is heavily used in microeconomics and company management, such as planning, product, transportation, technology and other issues, either to maximize the income or minimize the costs of a production scheme.

UNIT – VI: PROBABILITY

CHAPTER-13 PROBABILITY

Topic-1

Conditional Probability and Multiplication Theorem on Probability

Concepts Covered • Conditional Probability,
• Multiplication Theorem of Probability



Revision Notes

1. Basic Definition of Probability:

Let S and E be the **sample space** and **event** in an experiment respectively.



Key Words

Sample Space: A set in which all of the possible outcomes of a statistical experiment are represented as points.

Event: Event is a subset of a sample space. e.g.: Event of getting odd outcome in a throw of a die.

Then, Probability

$$= \frac{\text{Number of Favourable Events}}{\text{Total number of Elementary Events}} = \frac{n(E)}{n(S)}$$

$$0 \leq n(E) \leq n(S)$$

$$0 \leq P(E) \leq 1$$

Hence, if $P(E)$ denotes the probability of occurrence of an event E , then $0 \leq P(E) \leq 1$ and $P(\bar{E}) = 1 - P(E)$

such that $P(\bar{E})$ denotes the probability of non-occurrence of the event E .

⇒ Note that $P(\bar{E})$ can also be represented as $P(E')$.

2. Mutually Exclusive Or Disjoint Events:

Two events A and B are said to be mutually exclusive if occurrence of one prevents the occurrence of the other i.e., they can't occur simultaneously.

In this case, sets A and B are disjoint i.e., $A \cap B = \phi$.

3. Independent Events:

Two events are independent if the occurrence of one does not affect the occurrence of the other.

4. Exhaustive Events:

Two or more events say A , B and C of an experiment are said to be exhaustive events, if

(a) their union is the total sample space

$$\text{i.e., } A \cup B \cup C = S$$

(b) the event A , B and C are disjoint in pairs

i.e., $A \cap B = \phi$, $B \cap C = \phi$ and $C \cap A = \phi$.

(c) $P(A) + P(B) + P(C) = 1$.

➤ If A and B are mutually exhaustive events, then we always have

$$P(A \cap B) = 0 \quad [\text{As } n(A \cap B) = n(\phi) = 0]$$

$$\therefore P(A \cup B) = P(A) + P(B).$$

➤ If A , B and C are mutually exhaustive events, then we always have

$$P(A \cup B \cup C) = P(A) + P(B) + P(C)$$



Mnemonics

Concept: Independent and Mutually exclusive events.

I is not **ME**

ME is not **I**

Here, **I**: Independent Events

ME: Mutually Exclusive events

5. Conditional Probability:

By the conditional probability, we mean the probability of occurrence of event A when B has already

occurred.

The 'conditional probability of occurrence of event A when B has already occurred' is sometimes also called as probability of occurrence of event A w.r.t. B .

➤ $P(A|B) = \frac{P(A \cap B)}{P(B)}$, $B \neq \phi$ i.e., $P(B) \neq 0$

➤ $P(B|A) = \frac{P(A \cap B)}{P(A)}$, $A \neq \phi$ i.e., $P(A) \neq 0$

➤ $P(\bar{A}|B) = \frac{P(\bar{A} \cap B)}{P(B)}$, $P(B) \neq 0$

➤ $P(A|\bar{B}) = \frac{P(A \cap \bar{B})}{P(\bar{B})}$, $P(\bar{B}) \neq 0$

➤ $P(\bar{A}|\bar{B}) = \frac{P(\bar{A} \cap \bar{B})}{P(\bar{B})}$, $P(\bar{B}) \neq 0$

➤ $P(A|B) + P(\bar{A}|B) = 1$, $B \neq \phi$.



Key Fact

- Probability is originated from a gambler's dispute in 1654 concerning the division of a stake between two players whose game was interrupted before it close.



Key Formulae

(a) $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ i.e., $P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$

(b) $P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(C \cap A) + P(A \cap B \cap C)$

(c) $P(\bar{A} \cap B) = P(\text{only } B) = P(B - A) = P(B \text{ but not } A) = P(B) - P(A \cap B)$

(d) $P(A \cap \bar{B}) = P(\text{only } A) = P(A - B) = P(A \text{ but not } B) = P(A) - P(A \cap B)$

(e) $P(\bar{A} \cap \bar{B}) = P(\text{neither } A \text{ nor } B) = 1 - P(A \cup B)$

NOTE : EVENTS AND SYMBOLIC REPRESENTATIONS:

Verbal description of the event	Equivalent set notation
Event A	A
Not A	\bar{A} or A'
A or B (occurrence of atleast one A or B)	$A \cup B$ or $A + B$
A and B (simultaneous occurrence of both A and B)	$A \cap B$ or AB
A but not B (A occurs but B does not)	$A \cap \bar{B}$ or $A - B$
Neither A nor B	$\bar{A} \cap \bar{B}$
Atleast one A , B or C	$A \cup B \cup C$
All the three A , B and C	$A \cap B \cap C$



Key Facts

- The probability of living 110 years or more is about 1 in 7 million.

- If you are in the group of 23 people, there is a 50% chance that 2 of them share a birthday. If you are in a group of 70 people, that probability jumps to over 99%.

Topic-2**Bayes' Theorem****Concept Covered** • Bayes' Theorem**Revision Notes****BAYES' THEOREM:**

If $E_1, E_2, E_3, \dots, E_n$ are n non empty events constituting a partition of sample space S i.e., $E_1, E_2, E_3, \dots, E_n$ are pair wise disjoint and $E_1 \cup E_2 \cup E_3 \cup \dots \cup E_n = S$ and A is any event of non-zero probability, then

$$P(E_i|A) = \frac{P(E_i).P(A|E_i)}{\sum_{j=1}^n P(E_j).P(A|E_j)}, i = 1, 2, 3, \dots, n$$

For example, $P(E_1|A)$

$$= \frac{P(E_1).P(A|E_1)}{P(E_1).P(A|E_1) + P(E_2).P(A|E_2) + P(E_3).P(A|E_3)}$$

- Bayes' theorem is also known as the formula for the probability of causes.

- If $E_1, E_2, E_3, \dots, E_n$ form a partition of S and A be any event, then

$$P(A) = P(E_1).P(A|E_1) + P(E_2).P(A|E_2) + \dots + P(E_n).P(A|E_n)$$

$$[\because P(E_i \cap A) = P(E_i).P(A|E_i)]$$

- The probabilities $P(E_1), P(E_2), \dots, P(E_n)$ which are known before the experiment takes place are called **prior probabilities** and $P(A|E_n)$ are called **posterior probabilities**.

Topic-3**Random Variable and its Probability Distributions****Concepts Covered** • Random Variable, • Probability Distribution**Revision Notes****1. RANDOM VARIABLE:**

A random variable is a real valued function defined over the sample space of an experiment. In other words, a random variable is a real-valued function whose domain is the sample space of a random experiment. A random variable is usually denoted by uppercase letters X, Y, Z etc.

2. PROBABILITY DISTRIBUTION OF A RANDOM VARIABLE:

If the values of a random variable together with the corresponding probabilities are given, then this description is called a probability distribution of the random variable.

**Key Terms**

- **Discrete random variable:** It is a random variable which can take only finite or countable infinite number of values.
- **Continuous random variable:** A variable which can take any value between two given limits is called a continuous random variable.

**Key Formulae**

- Mean or Expectation of a random variable $X = \mu = \sum_{i=1}^n x_i P_i$